

H3K18ac polyclonal antibody
Purified Rabbit Polyclonal Antibody
Catalog # ABV11639**Specification**

H3K18ac polyclonal antibody - Product Information

Application	DB, E, WB
Primary Accession	P68431
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	15404

H3K18ac polyclonal antibody - Additional Information**Gene ID** 8350;8351;8352;8353;8354;8355;8356;8357;8358;8968**Other Names**

Histone 3

Target/Specificity

H3K18ac

Formulation

In PBS with 0.05% (W/V) sodium azide.

Handling

The antibody solution should be gently mixed before use.

Background Descriptions**Precautions**

H3K18ac polyclonal antibody is for research use only and not for use in diagnostic or therapeutic procedures.

H3K18ac polyclonal antibody - Protein Information**Name** H3C1 ([HGNC:4766](#))**Synonyms** H3FA, HIST1H3A**Function**

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.


Cellular Location
Nucleus. Chromosome.

H3K18ac polyclonal antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


H3K18ac polyclonal antibody - Images



A Dot Blot analysis was performed to test the cross reactivity of the antibody with peptides containing other modifications of histone H3 and an unmodified H3 sequence. 100 to 0.2 pmol of the peptide containing the respective histone modification were spotted on a membrane. The Fig shows a high specificity of the antibody for the modification of interest.



To determine the titer, an ELISA was performed using a serial dilution of the antibody. The antigen used was a peptide containing the histone modification of interest. By plotting the absorbance against the antibody dilution the titer of the antibody was estimated to be 1:8800.



HeLa cells treated with butyrate and histone extracts (15ug) were analysed by WB blot using the antibody.

H3K18ac polyclonal antibody - Background

Histones are the main constituents of the protein part of chromosomes of eukaryotic cells. They are rich in the amino acids arginine and lysine and have been greatly conserved during evolution. Histones pack the DNA into tight masses of chromatin. Two core histones of each class H2A, H2B, H3 and H4 assemble and are wrapped by 146 base pairs of DNA to form one octameric nucleosome. Histone tails undergo numerous post-translational modifications, which either directly or indirectly alter chromatin structure to facilitate transcriptional activation or repression or other nuclear processes. In addition to the genetic code, combinations of the different histone modifications reveal the so-called "histone code". Histone methylation and demethylation is dynamically regulated by respectively histone methyl transferases and histone demethylases. Acetylation of histone H3 is associated with active gene transcription.