

Anti-Collagen-I Antibody

Rabbit Polyclonal Antibody Catalog # ABV11815

Specification

Anti-Collagen-I Antibody - Product Information

Application DB
Primary Accession P02452
Reactivity Human
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 138911

Anti-Collagen-I Antibody - Additional Information

Gene ID 1277

Positive Control
Application & Usage

WB: 1-4 μg, Dot blot: 1-4 μg

Alias Symbol COL1A1

Other Names

Alpha-1 type I collagen, Collagen alpha-1(I) chain

AppearanceColorless liquid

Formulation

In PBS pH 7.2, 0.01 % BSA, 0.03 % ProClin® and 50 % glycerol

Reconstitution & Storage -20 °C

Background Descriptions

Precautions

Anti-Collagen-I Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Collagen-I Antibody - Protein Information

Name COL1A1

Function

Type I collagen is a member of group I collagen (fibrillar forming collagen).

Cellular Location



Secreted, extracellular space, extracellular matrix {ECO:0000255|PROSITE-ProRule:PRU00793}

Tissue Location


Forms the fibrils of tendon, ligaments and bones. In bones the fibrils are mineralized with calcium hydroxyapatite

Anti-Collagen-I Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Collagen-I Antibody - Images



Dot blot analysis of collagen using anti-Collagen antibody: Lane1 (rh Collagen-1); Lane2 (BSA)

Anti-Collagen-I Antibody - Background

Collagen, a major component of the extracellular matrix, is a fibrous protein that provides tensile strength to tissues giving them structural integrity. Collagen and its derivative, gelatin, have been widely used in medical, pharmaceutical and consumer products for more than 100 years. Collagens are fibrous, extracellular matrix proteins with high tensile strength and are the major components of connective tissue, such as tendons and cartilage. All collagens contain a triple helix domain and frequently show lateral self-association in order to form complex connective tissues. Several collagens also play a role in cell adhesion, important for maintaining normal tissue architecture and function. The extensive family of COL gene products (collagens) is composed of several chain types, including fibril-forming interstitial collagens (types I, II, III and V) and basement membrane collagens (type IV), each type containing multiple isoforms.