

Phospho Alpha-synuclein (Tyr125) Antibody

Catalog # ABV11950

Specification

Phospho Alpha-synuclein (Tyr125) Antibody - Product Information

Application WB, IHC Primary Accession P37840

Reactivity Human, Mouse, Rat

Host Rabbit Isotype Rabbit IgG Calculated MW 14460

Phospho Alpha-synuclein (Tyr125) Antibody - Additional Information

Gene ID 6622

Positive Control WB; HeLa, mouse brain, rat brain lysate,

IHC; human brain tissue

Application & Usage WB; 1:500 - 1:2000, IHC; 1:50 - 1:200

Other Names

NACP; PARK1; Alpha-synuclein; Non-A beta component of AD amyloid; Non-A4 component of

amyloid precursor; NACP

Target/Specificity

SNCA

Antibody Form

Liquid

Appearance

Colorless liquid

Handling

The antibody solution should be gently mixed before use

Reconstitution & Storage

-20°C

Background Descriptions

Precautions

Phospho Alpha-synuclein (Tyr125) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho Alpha-synuclein (Tyr125) Antibody - Protein Information

Name SNCA



Synonyms NACP, PARK1

Function

Neuronal protein that plays several roles in synaptic activity such as regulation of synaptic vesicle trafficking and subsequent neurotransmitter release (PubMed:28288128, PubMed:30404828, PubMed:20798282, PubMed:26442590). Participates as a monomer in synaptic vesicle exocytosis by enhancing vesicle priming, fusion and dilation of exocytotic fusion pores (PubMed:28288128, PubMed:30404828). Mechanistically, acts by increasing local Ca(2+) release from microdomains which is essential for the enhancement of ATP-induced exocytosis (PubMed: 30404828). Acts also as a molecular chaperone in its multimeric membrane-bound state, assisting in the folding of synaptic fusion components called SNAREs (Soluble NSF Attachment Protein REceptors) at presynaptic plasma membrane in conjunction with cysteine string protein-alpha/DNAIC5 (PubMed:20798282). This chaperone activity is important to sustain normal SNARE-complex assembly during aging (PubMed:20798282). Also plays a role in the regulation of the dopamine neurotransmission by associating with the dopamine transporter (DAT1) and thereby modulating its activity (PubMed: 26442590).

Cellular Location

Cytoplasm. Membrane. Nucleus. Synapse Secreted. Cell projection, axon {ECO:0000250|UniProtKB:O55042}. Note=Membrane-bound in dopaminergic neurons (PubMed:15282274). Expressed and colocalized with SEPTIN4 in dopaminergic axon terminals, especially at the varicosities (By similarity). {ECO:0000250|UniProtKB:O55042, ECO:0000269|PubMed:15282274}

Tissue Location

Highly expressed in presynaptic terminals in the central nervous system. Expressed principally in brain


Phospho Alpha-synuclein (Tyr125) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Phospho Alpha-synuclein (Tyr125) Antibody - Images





WB analysis of Alphasynuclein (pY125) expression in HeLa (A); mouse brain (B); rat brain (C) whole cell lysates



Immunohistochemical analysis of Alpha-synuclein (pY125) staining in human brain formalin fixed paraffin embedded tissue section.