

Goat Anti-Calretinin Antibody
Peptide-affinity purified goat antibody
Catalog # AF1182a**Specification**

Goat Anti-Calretinin Antibody - Product Information

Application	WB
Primary Accession	P07355
Other Accession	NP_001129487 , 302 , 12306 (mouse) , 56611 (rat)
Reactivity	Human, Mouse
Predicted	Rat, Pig, Cow
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	38604

Goat Anti-Calretinin Antibody - Additional Information**Gene ID** 302**Other Names**

Annexin A2, Annexin II, Annexin-2, Calpactin I heavy chain, Calpactin-1 heavy chain, Chromobindin-8, Lipocortin II, Placental anticoagulant protein IV, PAP-IV, Protein I, p36, ANXA2, ANX2, ANX2L4, CAL1H, LPC2D

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-Calretinin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-Calretinin Antibody - Protein Information**Name** ANXA2**Synonyms** ANX2, ANX2L4, CAL1H, LPC2D**Function**

Calcium-regulated membrane-binding protein whose affinity for calcium is greatly enhanced by

anionic phospholipids. It binds two calcium ions with high affinity. May be involved in heat-stress response. Inhibits PCSK9-enhanced LDLR degradation, probably reduces PCSK9 protein levels via a translational mechanism but also competes with LDLR for binding with PCSK9 (PubMed:18799458, PubMed:24808179, PubMed:22848640).

Cellular Location


Secreted, extracellular space, extracellular matrix, basement membrane. Melanosome. Note=In the lamina beneath the plasma membrane. Identified by mass spectrometry in melanosome fractions from stage I to stage IV. Translocated from the cytoplasm to the cell surface through a Golgi-independent mechanism

Goat Anti-Calretinin Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-Calretinin Antibody - Images



AF1182a staining (0.5 µg/ml) of A431 lysate (RIPA buffer, 35 µg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.

Goat Anti-Calretinin Antibody - Background

This gene encodes a member of the annexin family. Members of this calcium-dependent phospholipid-binding protein family play a role in the regulation of cellular growth and in signal transduction pathways. This protein functions as an autocrine factor which heightens osteoclast formation and bone resorption. This gene has three pseudogenes located on chromosomes 4, 9 and

10, respectively. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Goat Anti-Calretinin Antibody - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.

A role for the annexin A2 amino-terminal peptide in the plasmin-induced activation of human peripheral monocytes. Li Q, et al. Mol Immunol, 2010 Aug. PMID 20627396.

[A study of the single nucleotide polymorphism in seven genes (GHR, IGFBP3, IGFR1, IRS1, FMN1, ANXA2, TaGLN) in ethnic Russians and in patients with prostate cancer] Lisitskaia KV, et al. Mol Gen Mikrobiol Virusol, 2010. PMID 20540360.

Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614.

Role of annexin A2 in the production of infectious hepatitis C virus particles. Backes P, et al. J Virol, 2010 Jun. PMID 20335258.