

**Goat Anti-FXR1 Antibody**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF1450a****Specification**

---

**Goat Anti-FXR1 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P51114</a>
Other Accession	<a href="#">NP_001013457</a> , <a href="#">8087</a> , <a href="#">14359 (mouse)</a> , <a href="#">361927 (rat)</a>
Reactivity	Mouse
Predicted	Human, Rat, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5mg/ml
Isotype	IgG
Calculated MW	69721

**Goat Anti-FXR1 Antibody - Additional Information****Gene ID** 8087**Other Names**

Fragile X mental retardation syndrome-related protein 1, hFXR1p, FXR1

**Format**

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-FXR1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-FXR1 Antibody - Protein Information****Name** FXR1 {ECO:0000303|PubMed:7781595, ECO:0000312|HGNC:HGNC:4023}**Function**

mRNA-binding protein that acts as a regulator of mRNAs translation and/or stability, and which is required for various processes, such as neurogenesis, muscle development and spermatogenesis (PubMed:&lt;a href="http://www.uniprot.org/citations/17382880" target="\_blank"&gt;17382880&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/20417602" target="\_blank"&gt;20417602&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/30067974" target="\_blank"&gt;30067974&lt;/a&gt;),

PubMed:<a href="http://www.uniprot.org/citations/34731628" target="\_blank">34731628</a>, PubMed:<a href="http://www.uniprot.org/citations/35989368" target="\_blank">35989368</a>, PubMed:<a href="http://www.uniprot.org/citations/36306353" target="\_blank">36306353</a>). Specifically binds to AU-rich elements (AREs) in the 3'-UTR of target mRNAs (PubMed:<a href="http://www.uniprot.org/citations/17382880" target="\_blank">17382880</a>, PubMed:<a href="http://www.uniprot.org/citations/34731628" target="\_blank">34731628</a>). Promotes formation of some phase-separated membraneless compartment by undergoing liquid-liquid phase separation upon binding to AREs-containing mRNAs, leading to assemble mRNAs into cytoplasmic ribonucleoprotein granules that concentrate mRNAs with associated regulatory factors (By similarity). Required to activate translation of stored mRNAs during late spermatogenesis: acts by undergoing liquid-liquid phase separation to assemble target mRNAs into cytoplasmic ribonucleoprotein granules that recruit translation initiation factor EIF4G3 to activate translation of stored mRNAs in late spermatids (By similarity). Promotes translation of MYC transcripts by recruiting the eIF4F complex to the translation start site (PubMed:<a href="http://www.uniprot.org/citations/34731628" target="\_blank">34731628</a>). Acts as a negative regulator of inflammation in response to IL19 by promoting destabilization of pro-inflammatory transcripts (PubMed:<a href="http://www.uniprot.org/citations/30067974" target="\_blank">30067974</a>). Also acts as an inhibitor of inflammation by binding to TNF mRNA, decreasing TNF protein production (By similarity). Acts as a negative regulator of AMPA receptor GRIA2/GluA2 synthesis during long-lasting synaptic potentiation of hippocampal neurons by binding to GRIA2/GluA2 mRNA, thereby inhibiting its translation (By similarity). Regulates proliferation of adult neural stem cells by binding to CDKN1A mRNA and promoting its expression (By similarity). Acts as a regulator of sleep and synaptic homeostasis by regulating translation of transcripts in neurons (By similarity). Required for embryonic and postnatal development of muscle tissue by undergoing liquid-liquid phase separation to assemble target mRNAs into cytoplasmic ribonucleoprotein granules (PubMed:<a href="http://www.uniprot.org/citations/30770808" target="\_blank">30770808</a>). Involved in the nuclear pore complex localization to the nuclear envelope by preventing cytoplasmic aggregation of nucleoporins: acts by preventing ectopic phase separation of nucleoporins in the cytoplasm via a microtubule-dependent mechanism (PubMed:<a href="http://www.uniprot.org/citations/32706158" target="\_blank">32706158</a>).

### Cellular Location

Cytoplasm, Cytoplasmic ribonucleoprotein granule. Cytoplasm, Stress granule. Cytoplasm. Cell projection, dendrite {ECO:0000250|UniProtKB:Q61584}. Cell projection, dendritic spine {ECO:0000250|UniProtKB:Q61584}. Cell projection, axon {ECO:0000250|UniProtKB:Q61584}. Nucleus envelope. Postsynapse {ECO:0000250|UniProtKB:Q61584}. Note=Specifically localizes to cytoplasmic ribonucleoprotein membraneless compartments (By similarity). Localizes to stress granules following phosphorylation at Ser-420 by PAK1 (PubMed:20417602). Adjacent to Z-lines in muscles (By similarity). {ECO:0000250|UniProtKB:Q61584, ECO:0000269|PubMed:20417602}

### Tissue Location

Expressed in all tissues examined including heart, brain, kidney and testis (PubMed:7781595, PubMed:9259278). In brain, present at high level in neurons and especially in the Purkinje cells at the interface between the granular layer and the molecular layer (at protein level) (PubMed:9259278).

## Goat Anti-FXR1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Goat Anti-FXR1 Antibody - Images



AF1450a (0.2 µg/ml) staining of NIH/3T3 cell lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

#### Goat Anti-FXR1 Antibody - Background

The protein encoded by this gene is an RNA binding protein that interacts with the functionally-similar proteins FMR1 and FXR2. These proteins shuttle between the nucleus and cytoplasm and associate with polyribosomes, predominantly with the 60S ribosomal subunit. Three transcript variants encoding different isoforms have been found for this gene.

#### Goat Anti-FXR1 Antibody - References

Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. International Schizophrenia Consortium, et al. Nature, 2009 Aug 6. PMID 19571811.

Discrimination of common and unique RNA-binding activities among Fragile X mental retardation protein paralogs. Darnell JC, et al. Hum Mol Genet, 2009 Sep 1. PMID 19487368.

Alteration of expression of muscle specific isoforms of the fragile X related protein 1 (FXR1P) in facioscapulohumeral muscular dystrophy patients. Davidovic L, et al. J Med Genet, 2008 Oct. PMID 18628314.

Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348.

AU-rich-element-mediated upregulation of translation by FXR1 and Argonaute 2. Vasudevan S, et al. Cell, 2007 Mar 23. PMID 17382880.