

Goat Anti-GCDFP15 / PIP Antibody

Peptide-affinity purified goat antibody Catalog # AF1469a

Specification

Goat Anti-GCDFP15 / PIP Antibody - Product Information

Application IHC, WB Primary Accession P12273

Other Accession NP 002643, 5304

Reactivity
Human
Host
Clonality
Polyclonal
Concentration
100ug/200ul

Isotype IgG
Calculated MW 16572

Goat Anti-GCDFP15 / PIP Antibody - Additional Information

Gene ID 5304

Other Names

Prolactin-inducible protein, Gross cystic disease fluid protein 15, GCDFP-15, Prolactin-induced protein, Secretory actin-binding protein, SABP, gp17, PIP, GCDFP15, GPIP4

Format

0.5 mg lgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-GCDFP15 / PIP Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-GCDFP15 / PIP Antibody - Protein Information

Name PIP

Synonyms GCDFP15, GPIP4

Cellular Location

Secreted.

Tissue Location

Expressed in pathological conditions of the mammary gland and in several exocrine tissues, such




as the lacrimal, salivary, and sweat glands

Goat Anti-GCDFP15 / PIP Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Goat Anti-GCDFP15 / PIP Antibody - Images



AF1469a (2.5 μ g/ml) staining of paraffin embedded Human Breast. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



AF1469a (0.3 μ g/ml) staining of A431 cell lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-GCDFP15 / PIP Antibody - References

CLC- a novel susceptibility gene for allergic rhinitis? Bryborn M, et al. Allergy, 2010 Feb. PMID 19650845.





Tel: 858.875.1900 Fax: 858.875.1999

GCDFP-15 positive and TTF-1 negative primary lung neoplasms: a tissue microarray study of 381 primary lung tumors. Wang LJ, et al. Appl Immunohistochem Mol Morphol, 2009 Dec. PMID 19620839.

Defining the human deubiquitinating enzyme interaction landscape. Sowa ME, et al. Cell, 2009 Jul 23. PMID 19615732.

A functional and regulatory network associated with PIP expression in human breast cancer. Debily MA, et al. PLoS One, 2009. PMID 19262752.

Diagnostic utility of mammaglobin and GCDFP-15 in the identification of metastatic breast carcinoma in fluid specimens. Yan Z, et al. Diagn Cytopathol, 2009 Jul. PMID 19217055.