

Goat Anti-GPX3 Antibody
Peptide-affinity purified goat antibody
Catalog # AF1500a**Specification**

Goat Anti-GPX3 Antibody - Product Information

Application	WB, IHC
Primary Accession	P22352
Other Accession	NP_002075 , 2878 , 14778 (mouse) , 64317 (rat)
Reactivity	Mouse, Pig
Predicted	Human, Rat, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	25552

Goat Anti-GPX3 Antibody - Additional Information**Gene ID** 2878**Other Names**

Glutathione peroxidase 3, GPx-3, GSHPx-3, 1.11.1.9, Extracellular glutathione peroxidase, Plasma glutathione peroxidase, GPx-P, GSHPx-P, GPX3, GPXP

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-GPX3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-GPX3 Antibody - Protein Information**Name** GPX3 ([HGNC:4555](#))**Synonyms** GPXP**Function**

Protects cells and enzymes from oxidative damage, by catalyzing the reduction of hydrogen peroxide, lipid peroxides and organic hydroperoxide, by glutathione.

Cellular Location
Secreted.

Tissue Location
Secreted in plasma.

Goat Anti-GPX3 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-GPX3 Antibody - Images




AF1500a (1 µg/ml) staining of Mouse Heart lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.




AF1500a (20 µg/ml) staining of PFA-perfused cryosection of Porcine Kidney. Microwave antigen retrieval with citrate buffer pH 3, CY3-staining. Data obtained from Dr. Hrvoje Brzica, University of Zagreb, Croatia



EB07685 (2 μ g/ml) staining of Mouse Heart lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB07685 (4 μ g/ml) staining of paraffin embedded Human Kidney. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB07685 Negative Control showing staining of paraffin embedded Human Kidney, with no primary antibody.

Goat Anti-GPX3 Antibody - Background

This gene product belongs to the glutathione peroxidase family, which functions in the detoxification of hydrogen peroxide. It contains a selenocysteine (Sec) residue at its active site. The selenocysteine is encoded by the UGA codon, which normally signals translation termination. The 3' UTR of Sec-containing genes have a common stem-loop structure, the sec insertion sequence (SECIS), which is necessary for the recognition of UGA as a Sec codon rather than as a stop signal.

Goat Anti-GPX3 Antibody - References

Assessing oxidative pathway genes as risk factors for bipolar disorder. Fullerton JM, et al. Bipolar Disord, 2010 Aug. PMID 20712757.

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.

Functional glutathione peroxidase 3 polymorphisms associated with increased risk of Taiwanese patients with gastric cancer. Wang JY, et al. Clin Chim Acta, 2010 Oct 9. PMID 20576521.

Common polymorphisms in ITGA2, PON1 and THBS2 are associated with coronary atherosclerosis in a candidate gene association study of the Chinese Han population. Wang Y, et al. J Hum Genet, 2010 Aug. PMID 20485444.

Differential expression and distribution of placental glutathione peroxidases 1, 3 and 4 in normal and preeclamptic pregnancy. Mistry HD, et al. Placenta, 2010 May. PMID 20303587.