

**Goat Anti-GRM7 Antibody**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF1512a****Specification**

---

**Goat Anti-GRM7 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q14831</a>
Other Accession	<a href="#">NP_870989</a> , <a href="#">2917</a> , <a href="#">108073 (mouse)</a> , <a href="#">81672 (rat)</a>
Reactivity	Human
Predicted	Mouse, Rat
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	102251

**Goat Anti-GRM7 Antibody - Additional Information****Gene ID** 2917**Other Names**

Metabotropic glutamate receptor 7, mGluR7, GRM7, GPRC1G, MGLUR7

**Format**

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-GRM7 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-GRM7 Antibody - Protein Information****Name** GRM7**Synonyms** GPRC1G, MGLUR7**Function**

G-protein coupled receptor activated by glutamate that regulates axon outgrowth through the MAPK-cAMP-PKA signaling pathway during neuronal development (PubMed:&lt;a href="http://www.uniprot.org/citations/33500274" target="\_blank"&gt;33500274&lt;/a&gt;). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide- binding

proteins (G proteins) and modulates the activity of downstream effectors, such as adenylate cyclase that it inhibits (PubMed:<a href="http://www.uniprot.org/citations/9473604" target="\_blank">9473604</a>).

#### Cellular Location

Cell membrane; Multi-pass membrane protein

#### Tissue Location

Expressed in many areas of the brain, especially in the cerebral cortex, hippocampus, and cerebellum. Expression of GRM7 isoforms in non-neuronal tissues appears to be restricted to isoform 3 and isoform 4.

### Goat Anti-GRM7 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Goat Anti-GRM7 Antibody - Images



AF1512a (0.5 µg/ml) staining of Human Cerebellum lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

### Goat Anti-GRM7 Antibody - Background

L-glutamate is the major excitatory neurotransmitter in the central nervous system, and it activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors that have been divided into three groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5, and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3,

while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Multiple transcript variants encoding different isoforms have been found for this gene.

#### **Goat Anti-GRM7 Antibody - References**

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.

Comprehensive copy number variant (CNV) analysis of neuronal pathways genes in psychiatric disorders identifies rare variants within patients. Saus E, et al. J Psychiatr Res, 2010 Apr 14. PMID 20398908.

Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614.

Human variation in alcohol response is influenced by variation in neuronal signaling genes. Joslyn G, et al. Alcohol Clin Exp Res, 2010 May. PMID 20201926.

[Association between copy number variants within metabotropic glutamate receptors 7 gene and schizophrenia] Zhao YL, et al. Zhongguo Yi Xue Ke Xue Yuan Xue Bao, 2009 Dec. PMID 20078931.