10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Goat Anti-MAX Antibody

Peptide-affinity purified goat antibody

Catalog # AF1654a

Specification

Goat Anti-MAX Antibody - Product Information

Application	**IHC**
Primary Accession	P61244
Other Accession	NP_660087, 4149, 17187 (mouse)
Reactivity	**Human, Mouse, Rat**
Predicted	**Dog, Cow**
Host	**Goat**
Clonality	**Polyclonal**
Concentration	**100ug/200ul**
Isotype	**IgG**
Calculated MW	**18275**

Goat Anti-MAX Antibody - Additional Information

Gene ID 4149

Other Names
Protein max, Class D basic helix-loop-helix protein 4, bHLHd4, Myc-associated factor X, MAX, BHLHD4

Format
0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
Goat Anti-MAX Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-MAX Antibody - Protein Information

Name MAX (HGNC:6913)

Synonyms BHLHD4

Function
Transcription regulator. Forms a sequence-specific DNA- binding protein complex with MYC or MAD which recognizes the core sequence 5'-CAC[GA]TG-3'. The MYC:MAX complex is a transcriptional activator, whereas the MAD:MAX complex is a repressor. May repress transcription via the

recruitment of a chromatin remodeling complex containing H3 'Lys-9' histone methyltransferase activity. Represses MYC transcriptional activity from E-box elements.

Cellular Location
Nucleus. Cell projection, dendrite.

Tissue Location
High levels found in the brain, heart and lung while lower levels are seen in the liver, kidney and skeletal muscle

Goat Anti-MAX Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Goat Anti-MAX Antibody - Images



AF1654a (0.01 µg/ml) staining of Jurkat lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-MAX Antibody - Background

The protein encoded by this gene is a member of the basic helix-loop-helix leucine zipper (bHLHZ) family of transcription factors. It is able to form homodimers and heterodimers with other family members, which include Mad, Mxi1 and Myc. Myc is an oncoprotein implicated in cell proliferation, differentiation and apoptosis. The homodimers and heterodimers compete for a common DNA target site (the E box) and rearrangement among these dimer forms provides a complex system of transcriptional regulation. Multiple alternatively spliced transcript variants have been described for this gene but the full-length nature for some of them is unknown.

Goat Anti-MAX Antibody - References

Differentiation-associated miR-22 represses Max expression and inhibits cell cycle progression. Ting Y, et al. Biochem Biophys Res Commun, 2010 Apr 9. PMID 20214878.

Multiple loci influence erythrocyte phenotypes in the CHARGE Consortium. Ganesh SK, et al. Nat Genet, 2009 Nov. PMID 19862010.
Genome-wide association study of anthropometric traits in Korcula Island, Croatia. Polasek O, et al. Croat Med J, 2009 Feb. PMID 19260139.
Switch from Mnt-Max to Myc-Max induces p53 and cyclin D1 expression and apoptosis during cholestasis in mouse and human hepatocytes. Yang H, et al. Hepatology, 2009 Mar. PMID 19086036.
Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348.