

## **Goat Anti-NANOG Antibody**

Peptide-affinity purified goat antibody Catalog # AF1707b

### **Specification**

## **Goat Anti-NANOG Antibody - Product Information**

Application WB, IF Primary Accession Q9H9S0

Other Accession NP\_079141, 79923, 71950 (mouse), 414065

<u>(rat)</u>

Reactivity Human, Pig

Predicted Dog
Host Goat
Clonality Polyclonal
Concentration 100ug/200ul

Isotype IgG
Calculated MW 34620

### **Goat Anti-NANOG Antibody - Additional Information**

### **Gene ID** 79923

#### **Other Names**

Homeobox protein NANOG, Homeobox transcription factor Nanog, hNanog, NANOG

#### **Format**

0.5~mg lgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

Goat Anti-NANOG Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### **Goat Anti-NANOG Antibody - Protein Information**

### Name NANOG

#### **Function**

Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophectoderm lineages. Blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes. Acts



as a transcriptional activator or repressor. Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'-[CG][GA][CG]C[GC]ATTAN[GC]- 3'. Binds to the POU5F1/OCT4 promoter (PubMed:<a href="http://www.uniprot.org/citations/25825768" target="\_blank">25825768</a>). Able to autorepress its expression in differentiating (ES) cells: binds to its own promoter following interaction with ZNF281/ZFP281, leading to recruitment of the NuRD complex and subsequent repression of expression. When overexpressed, promotes cells to enter into S phase and proliferation.

# **Cellular Location**

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108, ECO:0000269|PubMed:15983365}

#### **Tissue Location**


Expressed in testicular carcinoma and derived germ cell tumors (at protein level). Expressed in fetal gonads, ovary and testis. Also expressed in ovary teratocarcinoma cell line and testicular embryonic carcinoma. Not expressed in many somatic organs and oocytes.

## **Goat Anti-NANOG Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### **Goat Anti-NANOG Antibody - Images**



AF1707b (0.03μg/ml) staining of Human Ovary lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.





AF1707b (5ug/ml) staining (green) parts of a colony of induced pluriform stem cells derived from Human Keratinocytes. Data kindly provided by CMRB, Center of Regenerative Medicine in Barcelona, Spain.

# **Goat Anti-NANOG Antibody - References**

Hedgehog controls neural stem cells through p53-independent regulation of Nanog. Po A, et al. EMBO J, 2010 Aug 4. PMID 20581804.

NANOG regulates glioma stem cells and is essential in vivo acting in a cross-functional network with GLI1 and p53. Zbinden M, et al. EMBO J, 2010 Aug 4. PMID 20581802.

A distinct expression pattern in mammalian testes indicates a conserved role for NANOG in spermatogenesis. Kuijk EW, et al. PLoS One, 2010 Jun 7. PMID 20539761.

Expression profile of the embryonic markers nanog, OCT-4, SSEA-1, SSEA-4, and frizzled-9 receptor in human periodontal ligament mesenchymal stem cells. Trubiani O, et al. J Cell Physiol, 2010 Oct. PMID 20458727.

Novel candidate cancer genes identified by a large-scale cross-species comparative oncogenomics approach. Mattison J, et al. Cancer Res, 2010 Feb 1. PMID 20103622.