

Goat Anti-ORP11 / OSBPL11 Antibody
Peptide-affinity purified goat antibody
Catalog # AF1759a**Specification**

Goat Anti-ORP11 / OSBPL11 Antibody - Product Information

Application	WB
Primary Accession	Q9BXB4
Other Accession	NP_073613 , 114885
Reactivity	Human
Predicted	Mouse, Rat, Dog, Cow
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	83643

Goat Anti-ORP11 / OSBPL11 Antibody - Additional Information**Gene ID** 114885**Other Names**

Oxysterol-binding protein-related protein 11, ORP-11, OSBP-related protein 11, OSBPL11, ORP11, OSBP12

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-ORP11 / OSBPL11 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-ORP11 / OSBPL11 Antibody - Protein Information**Name** OSBPL11**Synonyms** ORP11, OSBP12**Function**

Plays a role in regulating ADIPOQ and FABP4 levels in differentiating adipocytes and is also involved in regulation of adipocyte triglyceride storage (PubMed:23028956). Weakly binds

25-hydroxycholesterol (PubMed:17428193).

Cellular Location

Late endosome membrane. Golgi apparatus, trans-Golgi network membrane. Note=Localizes at the Golgi- late endosome interface

Tissue Location


Present at highest levels in ovary, testis, kidney, liver, stomach, brain, and adipose tissue. Strong expression (at protein level) in epithelial cells of kidney tubules, testicular tubules, caecum, and skin (PubMed:20599956). Present at low levels in subcutaneous and visceral adipose tissue (at protein level)(PubMed:23028956).

Goat Anti-ORP11 / OSBPL11 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-ORP11 / OSBPL11 Antibody - Images



AF1759a staining (1 µg/ml) of Human Brain lysate (RIPA buffer, 30 µg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.

Goat Anti-ORP11 / OSBPL11 Antibody - Background

This gene encodes a member of the oxysterol-binding protein (OSBP) family, a group of intracellular lipid receptors. Like most members, the encoded protein contains an N-terminal pleckstrin homology domain and a highly conserved C-terminal OSBP-like sterol-binding domain.

Goat Anti-ORP11 / OSBPL11 Antibody - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.

Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121.

Defining the human deubiquitinating enzyme interaction landscape. Sowa ME, et al. Cell, 2009 Jul 23. PMID 19615732.

Association of OSBPL11 gene polymorphisms with cardiovascular disease risk factors in obesity. Bouchard L, et al. Obesity (Silver Spring), 2009 Jul. PMID 19325544.

Global, in vivo, and site-specific phosphorylation dynamics in signaling networks. Olsen JV, et al. Cell, 2006 Nov 3. PMID 17081983.