

## **Goat Anti-ORP2 Antibody**

Peptide-affinity purified goat antibody Catalog # AF1760a

### **Specification**

## **Goat Anti-ORP2 Antibody - Product Information**

Application WB
Primary Accession O9H1P3

Other Accession <u>NP\_653081</u>, <u>9885</u>, <u>228983 (mouse)</u>

Reactivity
Predicted
Host
Clonality
Concentration
Human
Mouse, Rat
Goat
Polyclonal
100ug/200ul

Isotype IgG Calculated MW 55201

## **Goat Anti-ORP2 Antibody - Additional Information**

### **Gene ID 9885**

## **Other Names**

Oxysterol-binding protein-related protein 2, ORP-2, OSBP-related protein 2, OSBPL2, KIAA0772, ORP2

#### **Format**

0.5~mg lgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

Goat Anti-ORP2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **Goat Anti-ORP2 Antibody - Protein Information**

### Name OSBPL2

Synonyms KIAA0772, ORP2

# **Function**

Intracellular transport protein that binds sterols and phospholipids and mediates lipid transport between intracellular compartments. Increases plasma membrane cholesterol levels and decreases phosphatidylinositol-4,5-bisphosphate levels in the cell membrane (PubMed:<a



href="http://www.uniprot.org/citations/30581148" target="\_blank">30581148</a>). Binds phosphoinositides, such as phosphatidylinositol-4,5-bisphosphate (PubMed:<a href="http://www.uniprot.org/citations/30581148" target="\_blank">30581148</a>). Exhibits strong binding to phosphatidic acid and weak binding to phosphatidylinositol 3-phosphate (PubMed:<a href="http://www.uniprot.org/citations/11279184" target="\_blank">11279184</a>). Binds cholesterol, dehydroergosterol, 22(R)-hydroxycholesterol and 25-hydroxycholesterol (in vitro) (PubMed:<a href="http://www.uniprot.org/citations/17428193" target="\_blank">17428193</a>, PubMed:<a href="http://www.uniprot.org/citations/19224871" target="\_blank">19224871</a><br/>, PubMed:<a href="http://www.uniprot.org/citations/30581148" target="\_blank">30581148</a>).

#### **Cellular Location**

Cytoplasm, cytosol. Lipid droplet. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Note=Detected on the surface of cytosolic lipid droplets (PubMed:19224871). Recruited to the cell membrane by phosphatidylinositol-phosphate binding (PubMed:30581148)

Tissue Location Widely expressed.

## **Goat Anti-ORP2 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### Goat Anti-ORP2 Antibody - Images



AF1760a staining (1.5  $\mu$ g/ml) of Human Heart lysate (RIPA buffer, 30  $\mu$ g total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.

### Goat Anti-ORP2 Antibody - Background





Tel: 858.875.1900 Fax: 858.875.1999

This gene encodes a member of the oxysterol-binding protein (OSBP) family, a group of intracellular lipid receptors. Most members contain an N-terminal pleckstrin homology domain and a highly conserved C-terminal OSBP-like sterol-binding domain, although some members contain only the sterol-binding domain. This encoded protein contains only the sterol-binding domain. In vitro studies have shown that the encoded protein can bind strongly to phosphatic acid and weakly to phosphatidylinositol 3-phosphate, but cannot bind to 25-hydroxycholesterol. The protein associates with the Golgi apparatus. Transcript variants encoding different isoforms have been described.

## Goat Anti-ORP2 Antibody - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.

Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121.

OSBPL10, a novel candidate gene for high triglyceride trait in dyslipidemic Finnish subjects, regulates cellular lipid metabolism. Perttil∏ J, et al. J Mol Med, 2009 Aug. PMID 19554302.

OSBP-related protein 2 is a sterol receptor on lipid droplets that regulates the metabolism of neutral lipids. Hynynen R, et al. | Lipid Res, 2009 Jul. PMID 19224871.

The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334.