

Goat Anti-PALS2 / VAM1 / MPP6 Antibody
Peptide-affinity purified goat antibody
Catalog # AF1781a**Specification**

Goat Anti-PALS2 / VAM1 / MPP6 Antibody - Product Information

Application	WB
Primary Accession	O9NZW5
Other Accession	NP_057531 , 51678 , 56524 (mouse)
Reactivity	Human
Predicted	Mouse, Rat, Pig, Dog, Cow
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	61117

Goat Anti-PALS2 / VAM1 / MPP6 Antibody - Additional Information**Gene ID** 51678**Other Names**

MAGUK p55 subfamily member 6, Veli-associated MAGUK 1, VAM-1, MPP6, VAM1

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-PALS2 / VAM1 / MPP6 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-PALS2 / VAM1 / MPP6 Antibody - Protein Information**Name** PALS2 ([HGNC:18167](#))**Synonyms** MPP6, VAM1**Cellular Location**

Membrane; Peripheral membrane protein

Tissue Location

Abundant in testis, brain, and kidney with lower levels detectable in other tissues

Goat Anti-PALS2 / VAM1 / MPP6 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-PALS2 / VAM1 / MPP6 Antibody - Images



AF1781a staining (0.3 µg/ml) of 293 lysate (RIPA buffer, 35 µg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.

Goat Anti-PALS2 / VAM1 / MPP6 Antibody - Background

Members of the peripheral membrane-associated guanylate kinase (MAGUK) family function in tumor suppression and receptor clustering by forming multiprotein complexes containing distinct sets of transmembrane, cytoskeletal, and cytoplasmic signaling proteins. All MAGUKs contain a PDZ-SH3-GUK core and are divided into 4 subfamilies, DLG-like (see DLG1; MIM 601014), ZO1-like (see TJP1; MIM 601009), p55-like (see MPP1; MIM 305360), and LIN2-like (see CASK; MIM 300172), based on their size and the presence of additional domains. MPP6 is a member of the p55-like MAGUK subfamily (Tseng et al., 2001 [PubMed 11311936]).

Goat Anti-PALS2 / VAM1 / MPP6 Antibody - References

Prospective 7 year follow up imaging study comparing radiography, ultrasonography, and magnetic resonance imaging in rheumatoid arthritis finger joints. Scheel AK, et al. Ann Rheum Dis, 2006 May. PMID 16192290.

Immunoaffinity profiling of tyrosine phosphorylation in cancer cells. Rush J, et al. Nat Biotechnol, 2005 Jan. PMID 15592455.

The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334.

A protein interaction framework for human mRNA degradation. Lehner B, et al. Genome Res, 2004 Jul. PMID 15231747.

Protein trafficking and anchoring complexes revealed by proteomic analysis of inward rectifier potassium channel (Kir2.x)-associated proteins. Leonoudakis D, et al. J Biol Chem, 2004 May 21.

PMID 15024025.