

Goat Anti-PRUNE2 / BMCC1 Antibody
Peptide-affinity purified goat antibody
Catalog # AF1872a**Specification**

Goat Anti-PRUNE2 / BMCC1 Antibody - Product Information

Application	WB
Primary Accession	Q8WUY3
Other Accession	NP_620173 , 158471
Reactivity	Human
Predicted	Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	340635

Goat Anti-PRUNE2 / BMCC1 Antibody - Additional Information**Gene ID** 158471**Other Names**

Protein prune homolog 2, BNIP2 motif-containing molecule at the C-terminal region 1, PRUNE2, BMCC1, BNIPXL, C9orf65, KIAA0367

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-PRUNE2 / BMCC1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-PRUNE2 / BMCC1 Antibody - Protein Information**Name** PRUNE2**Synonyms** BMCC1, BNIPXL, C9orf65, KIAA0367**Function**

May play an important role in regulating differentiation, survival and aggressiveness of the tumor cells.

Cellular Location

Cytoplasm.

Tissue Location

A high level of expression seen in the nervous system (brain, cerebellum and spinal cord) as well as adrenal gland. Expressed at high levels in noneuroblastoma, rhabdomyosarcoma, melanoma and some osteosarcoma cell lines, whereas at only low levels in cancer cell lines of liver, breast, thyroid and colon. Expression is significantly higher in favorable tumors than aggressive ones

Goat Anti-PRUNE2 / BMCC1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-PRUNE2 / BMCC1 Antibody - Images

AF1872a (1 µg/ml) staining of Human Brain lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.