10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Goat Anti-RASSF8 Antibody

Peptide-affinity purified goat antibody
Catalog # AF1918a

Specification

Goat Anti-RASSF8 Antibody - Product Information

Application	**WB**
Primary Accession	Q8NHQ8
Other Accession	NP_009142, 11228, 71323 (mouse)
Reactivity	**Human**
Predicted	**Mouse, Cow**
Host	**Goat**
Clonality	**Polyclonal**
Concentration	**100ug/200ul**
Isotype	**IgG**
Calculated MW	**48327**

Goat Anti-RASSF8 Antibody - Additional Information

Gene ID 11228

Other Names
Ras association domain-containing protein 8, Carcinoma-associated protein HOJ-1, RASSF8, C12orf2

Format
0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
Goat Anti-RASSF8 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-RASSF8 Antibody - Protein Information

Name RASSF8

Synonyms C12orf2

Tissue Location
Widely expressed as a 6.2 kb transcript. A 2.2 kb alternatively spliced transcript is expressed exclusively in testis

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Goat Anti-RASSF8 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Goat Anti-RASSF8 Antibody - Images



AF1918a (0.3 µg/ml) staining of Human Testis lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-RASSF8 Antibody - References

The RASSF8 candidate tumor suppressor inhibits cell growth and regulates the Wnt and NF-kappaB signaling pathways. Lock FE, et al. Oncogene, 2010 Jul 29. PMID 20514026.
Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614.
Common polymorphisms in D12S1034 flanking genes RASSF8 and BHLHB3 are not associated with lung adenocarcinoma risk. Falvella FS, et al. Lung Cancer, 2007 Apr. PMID 17194498.
Identification of RASSF8 as a candidate lung tumor suppressor gene. Falvella FS, et al. Oncogene, 2006 Jun 29. PMID 16462760.
Diversification of transcriptional modulation: large-scale identification and characterization of putative alternative promoters of human genes. Kimura K, et al. Genome Res, 2006 Jan. PMID 16344560.