

Goat Anti-SART1 Antibody

Peptide-affinity purified goat antibody Catalog # AF1959a

Specification

Goat Anti-SART1 Antibody - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Concentration Isotype Calculated MW WB O43290 NP_005137, 9092, 20227 (mouse), 29678 (rat) Human Mouse, Rat, Cow Goat Polyclonal 100ug/200ul IgG 90255

Goat Anti-SART1 Antibody - Additional Information

Gene ID 9092

Other Names

U4/U6.U5 tri-snRNP-associated protein 1, SNU66 homolog, hSnu66, Squamous cell carcinoma antigen recognized by T-cells 1, SART-1, hSART-1, U4/U6.U5 tri-snRNP-associated 110 kDa protein, Hom s 1, SART1

Format

0.5 mg lgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-SART1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-SART1 Antibody - Protein Information

Name SART1

Function

Plays a role in mRNA splicing as a component of the U4/U6-U5 tri-snRNP, one of the building blocks of the spliceosome. May also bind to DNA.

Cellular Location



Nucleus. Note=Found in the nucleus of mitogen- activated peripheral blood mononuclear cells (PBMCs), tumor cells, or normal cell lines, but not in normal tissues except testis and fetal liver or in unstimulated PBMCs, suggesting preferential expression in proliferating cells

Tissue Location Ubiquitously expressed.

Goat Anti-SART1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Goat Anti-SART1 Antibody - Images

-	250kDa 150kDa 100kDa 75kDa
	50kDa
	37kDa
	25kDa 20kDa

AF1959a (0.03 μ g/ml) staining of Hela lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-SART1 Antibody - Background

This gene encodes two proteins, the SART1(800) protein expressed in the nucleus of the majority of proliferating cells, and the SART1(259) protein expressed in the cytosol of epithelial cancers. The SART1(259) protein is translated by the mechanism of -1 frameshifting during posttranscriptional regulation; its full-length sequence is not published yet. The two encoded proteins are thought to be involved in the regulation of proliferation. Both proteins have tumor-rejection antigens. The SART1(259) protein possesses tumor epitopes capable of inducing HLA-A2402-restricted cytotoxic T lymphocytes in cancer patients. This SART1(259) antigen may be useful in specific immunotherapy for cancer patients and may serve as a paradigmatic tool for the diagnosis and treatment of patients with atopy. The SART1(259) protein is found to be essential for the recruitment of the tri-snRNP to the pre-spliceosome in the spliceosome assembly pathway.

Goat Anti-SART1 Antibody - References

Association of mitotic regulation pathway polymorphisms with pancreatic cancer risk and outcome. Couch FJ, et al. Cancer Epidemiol Biomarkers Prev, 2010 Jan. PMID 20056645.



Defining the human deubiquitinating enzyme interaction landscape. Sowa ME, et al. Cell, 2009 Jul 23. PMID 19615732.

Hypoxia-associated factor, a novel E3-ubiquitin ligase, binds and ubiquitinates hypoxia-inducible factor 1alpha, leading to its oxygen-independent degradation. Koh MY, et al. Mol Cell Biol, 2008 Dec. PMID 18838541.

Influence of multiple genetic polymorphisms on genitourinary morbidity after carbon ion radiotherapy for prostate cancer. Suga T, et al. Int J Radiat Oncol Biol Phys, 2008 Nov 1. PMID 18374504.

Systematic analysis of the protein interaction network for the human transcription machinery reveals the identity of the 7SK capping enzyme. Jeronimo C, et al. Mol Cell, 2007 Jul 20. PMID 17643375.