

Goat Anti-SEL1L Antibody

Peptide-affinity purified goat antibody Catalog # AF1966a

Specification

Goat Anti-SEL1L Antibody - Product Information

Application WB
Primary Accession Q9UBV2

Other Accession NP 005056, 6400, 20338 (mouse), 314352 (rat)

Reactivity Human

Predicted Mouse, Rat, Pig, Dog, Cow

Host Goat
Clonality Polyclonal
Concentration 100ug/200ul

Isotype IgG
Calculated MW 88755

Goat Anti-SEL1L Antibody - Additional Information

Gene ID 6400

Other Names

Protein sel-1 homolog 1, Suppressor of lin-12-like protein 1, Sel-1L, SEL1L, TSA305

Format

0.5 mg lgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-SEL1L Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-SEL1L Antibody - Protein Information

Name SEL1L

Synonyms TSA305

Function

Plays a role in the endoplasmic reticulum quality control (ERQC) system also called ER-associated degradation (ERAD) involved in ubiquitin-dependent degradation of misfolded endoplasmic reticulum proteins (PubMed:16186509, PubMed:<a href="http://www.uniprot.org/citations/29997207"



target="_blank">29997207). Enhances SYVN1 stability. Plays a role in LPL maturation and secretion. Required for normal differentiation of the pancreas epithelium, and for normal exocrine function and survival of pancreatic cells. May play a role in Notch signaling.

Cellular Location

Endoplasmic reticulum membrane; Single-pass type I membrane protein

Tissue Location

Highly expressed in pancreas.

Goat Anti-SEL1L Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Goat Anti-SEL1L Antibody - Images



AF1966a (1 μ g/ml) staining of HepG2 lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-SEL1L Antibody - References

Identification of type 2 diabetes-associated combination of SNPs using support vector machine. Ban HJ, et al. BMC Genet, 2010 Apr 23. PMID 20416077.

A luminal flavoprotein in endoplasmic reticulum-associated degradation. Riemer J, et al. Proc Natl Acad Sci U S A, 2009 Sep 1. PMID 19706418.

EDEM1 recognition and delivery of misfolded proteins to the SEL1L-containing ERAD complex. Cormier JH, et al. Mol Cell, 2009 Jun 12. PMID 19524542.

Functional characterization of two secreted SEL1L isoforms capable of exporting unassembled substrate. Cattaneo M, et al. J Biol Chem, 2009 Apr 24. PMID 19204006.

Cln6 mutants associated with neuronal ceroid lipofuscinosis are degraded in a proteasome-dependent manner. Oresic K, et al. Biosci Rep, 2009 Jun. PMID 18811591.