

**Goat Anti-Triosephosphate isomerase Antibody**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF2113a****Specification**

---

**Goat Anti-Triosephosphate isomerase Antibody - Product Information**

|                   |   |
|-------------------|---|
| Application       | WB, IHC   |
| Primary Accession | <a href="#">P60174</a>                              |
| Other Accession   | <a href="#">NP_001152759</a> , <a href="#">7167</a> |
| Reactivity        | Human, Mouse  |
| Predicted         | Rat, Dog, Cow                                       |
| Host              | Goat  |
| Clonality         | Polyclonal  |
| Concentration     | 0.5mg/ml  |
| Isotype           | IgG   |
| Calculated MW     | 26669   |

**Goat Anti-Triosephosphate isomerase Antibody - Additional Information****Gene ID** 7167**Other Names**

Triosephosphate isomerase, TIM, 5.3.1.1, Triose-phosphate isomerase, TPI1, TPI

**Format**

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-Triosephosphate isomerase Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-Triosephosphate isomerase Antibody - Protein Information****Name** TPI1**Synonyms** TPI**Function**

Triosephosphate isomerase is an extremely efficient metabolic enzyme that catalyzes the interconversion between dihydroxyacetone phosphate (DHAP) and D-glyceraldehyde-3-phosphate (G3P) in glycolysis and gluconeogenesis.

**Cellular Location**

Cytoplasm {ECO:0000255|PROSITE-ProRule:PRU10127}.


**Goat Anti-Triosephosphate isomerase Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Goat Anti-Triosephosphate isomerase Antibody - Images**

AF2113a staining (0.003 µg/ml) of human liver lysate (RIPA buffer, 30 µg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.



AF2113a (2 µg/ml) staining of paraffin embedded Human Liver Steamed antigen retrieval with Tris/EDTA buffer pH 9, HRP-staining.

**Goat Anti-Triosephosphate isomerase Antibody - Background**

This gene encodes an enzyme, consisting of two identical proteins, which catalyzes the isomerization of glyceraldehydes 3-phosphate (G3P) and dihydroxy-acetone phosphate (DHAP) in glycolysis and gluconeogenesis. Mutations in this gene are associated with triosephosphate isomerase deficiency. Pseudogenes have been identified on chromosomes 1, 4, 6 and 7. Alternative splicing results in multiple transcript variants.

#### **Goat Anti-Triosephosphate isomerase Antibody - References**

Proteome analysis of the thalamus and cerebrospinal fluid reveals glycolysis dysfunction and potential biomarkers candidates for schizophrenia. Martins-de-Souza D, et al. J Psychiatr Res, 2010 May 14. PMID 20471030. Triose phosphate isomerase deficiency associated with two novel mutations in TPI gene. Fermo E, et al. Eur J Haematol, 2010 Aug. PMID 20374271. [Identification and expression of two new secretory proteins associated with prostate cancer] Qian XL, et al. Yi Chuan, 2010 Mar. PMID 20233700. Proteome analysis of schizophrenia patients Wernicke's area reveals an energy metabolism dysregulation. Martins-de-Souza D, et al. BMC Psychiatry, 2009 Apr 30. PMID 19405953. Analysis of TPI gene promoter variation in three sub-Saharan Africa population samples. Manco L, et al. Am J Hum Biol, 2009 Jan-Feb. PMID 18792062.