

Goat Anti-VPS28 Antibody
Peptide-affinity purified goat antibody
Catalog # AF2151a

Specification

Goat Anti-VPS28 Antibody - Product Information

Application	WB, IHC
Primary Accession	Q9UK41
Other Accession	NP_057292 , 51160 , 66914 (mouse)
Reactivity	Human
Predicted	Mouse, Rat, Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	25425

Goat Anti-VPS28 Antibody - Additional Information

Gene ID 51160

Other Names

Vacuolar protein sorting-associated protein 28 homolog, H-Vps28, ESCRT-I complex subunit VPS28, VPS28

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-VPS28 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-VPS28 Antibody - Protein Information

Name VPS28

Function

Component of the ESCRT-I complex, a regulator of vesicular trafficking process.

Cellular Location

Cell membrane. Late endosome membrane; Peripheral membrane protein

Goat Anti-VPS28 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-VPS28 Antibody - Images



AF2151a staining (1 µg/ml) of Human Testis lysate (RIPA buffer, 35 µg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.



AF2151a (10 µg/ml) staining of paraffin embedded Human Testis. Microwaved antigen retrieval with Tris/EDTA buffer pH9, HRP-staining.

Goat Anti-VPS28 Antibody - Background

This gene encodes a protein involved in endosomal sorting of cell surface receptors via a

multivesicular body/late endosome pathway. The encoded protein is one of the three subunits of the ESCRT-I complex (endosomal complexes required for transport) involved in the sorting of ubiquitinated proteins. The two other subunits of ESCRT-I are vesicular protein sorting 23, also known as tumor susceptibility gene 101 (TSG101), and vesicular protein sorting 37. Two alternative transcripts encoding different isoforms have been described. Additional alternative transcripts may exist but the proteins encoded by these transcripts have not been verified experimentally.

Goat Anti-VPS28 Antibody - References

Budding of filamentous and non-filamentous influenza A virus occurs via a VPS4 and VPS28-independent pathway. Bruce EA, et al. Virology, 2009 Aug 1. PMID 19524996.
An empirical framework for binary interactome mapping. Venkatesan K, et al. Nat Methods, 2009 Jan. PMID 19060904.
Regulation of Tsg101 expression by the steadiness box: a role of Tsg101-associated ligase. McDonald B, et al. Mol Biol Cell, 2008 Feb. PMID 18077552.
Towards a proteome-scale map of the human protein-protein interaction network. Rual JF, et al. Nature, 2005 Oct 20. PMID 16189514.
A human protein-protein interaction network: a resource for annotating the proteome. Stelzl U, et al. Cell, 2005 Sep 23. PMID 16169070.