

Goat Anti-VPS37C Antibody

Peptide-affinity purified goat antibody Catalog # AF2154a

Specification

Goat Anti-VPS37C Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Concentration Isotype Calculated MW WB, IHC <u>A5D8V6</u> <u>NP_060436</u>, <u>55048</u> Human Goat Polyclonal 100ug/200ul IgG 38659

Goat Anti-VPS37C Antibody - Additional Information

Gene ID 55048

Other Names Vacuolar protein sorting-associated protein 37C, hVps37C, ESCRT-I complex subunit VPS37C, VPS37C, PML39

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-VPS37C Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-VPS37C Antibody - Protein Information

Name VPS37C

Synonyms PML39

Function

Component of the ESCRT-I complex, a regulator of vesicular trafficking process. Required for the sorting of endocytic ubiquitinated cargos into multivesicular bodies. May be involved in cell growth and differentiation.



Cellular Location


Late endosome membrane; Peripheral membrane protein. Note=Probably associates with membranes Recruited to the plasma membrane by HIV-1

Goat Anti-VPS37C Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Goat Anti-VPS37C Antibody - Images



AF2154a (0.03g/ml) staining of human duodenum lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



AF2154a (3.8 μ g/ml) staining of paraffin embedded Human Prostate. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

Goat Anti-VPS37C Antibody - Background



VPS37C is a subunit of ESCRT-I (endosomal sorting complex required for transport I), a complex in the class E vacuolar protein sorting (VPS) pathway required for sorting ubiquitinated transmembrane proteins into internal vesicles of multivesicular bodies (Eastman et al., 2005 [PubMed 15509564]).

Goat Anti-VPS37C Antibody - References

Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614.

Association between genetic variants in VEGF, ERCC3 and occupational benzene haematotoxicity. Hosgood HD 3rd, et al. Occup Environ Med, 2009 Dec. PMID 19773279.

Identification of human VPS37C, a component of endosomal sorting complex required for transport-I important for viral budding. Eastman SW, et al. J Biol Chem, 2005 Jan 7. PMID 15509564.

Complete sequencing and characterization of 21,243 full-length human cDNAs. Ota T, et al. Nat Genet, 2004 Jan. PMID 14702039.

Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. Strausberg RL, et al. Proc Natl Acad Sci U S A, 2002 Dec 24. PMID 12477932.