

Goat Anti-HTR7 / 5-HT7 Antibody
Peptide-affinity purified goat antibody
Catalog # AF2182a**Specification**

Goat Anti-HTR7 / 5-HT7 Antibody - Product Information

Application	WB
Primary Accession	P34969
Other Accession	NP_000863 , 3363
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	53555

Goat Anti-HTR7 / 5-HT7 Antibody - Additional Information**Gene ID** 3363**Other Names**

5-hydroxytryptamine receptor 7, 5-HT-7, 5-HT7, 5-HT-X, Serotonin receptor 7, HTR7

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-HTR7 / 5-HT7 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-HTR7 / 5-HT7 Antibody - Protein Information**Name** HTR7**Function**

This is one of the several different receptors for 5- hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone, and a mitogen. The activity of this receptor is mediated by G proteins that stimulate adenylate cyclase.

Cellular Location

Cell membrane; Multi-pass membrane protein.

Tissue Location

Isoform A is the predominant isoform in spleen, caudate and hippocampus. Isoform B is expressed at lower levels Isoform D is a minor isoform in terms of expression

Goat Anti-HTR7 / 5-HT7 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-HTR7 / 5-HT7 Antibody - Images



AF2182a (1 µg/ml) staining of KELLY lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-HTR7 / 5-HT7 Antibody - Background

The neurotransmitter, serotonin, is thought to play a role in various cognitive and behavioral functions. The serotonin receptor encoded by this gene belongs to the superfamily of G protein-coupled receptors and the gene is a candidate locus for involvement in autistic disorder and other neuropsychiatric disorders. Three splice variants have been identified which encode proteins that differ in the length of their carboxy terminal ends.

Goat Anti-HTR7 / 5-HT7 Antibody - References

Physiogenomic analysis of statin-treated patients: domain-specific counter effects within the ACACB gene on low-density lipoprotein cholesterol? Ruaño G, et al. Pharmacogenomics, 2010 Jul. PMID 20602615.

Association study of 182 candidate genes in anorexia nervosa. Pinheiro AP, et al. Am J Med Genet B Neuropsychiatr Genet, 2010 Jul. PMID 20468064.

Association study of the serotoninergic system in migraine in the Spanish population. Corominas R, et al. Am J Med Genet B Neuropsychiatr Genet, 2010 Jan 5. PMID 19455600.

Expression of 5-HT1A and 5-HT7 receptors in Caco-2 cells and their role in the regulation of serotonin transporter activity. Iceta R, et al. J Physiol Pharmacol, 2009 Mar. PMID 19439818.

Association analysis of serotonin receptor 7 gene (HTR7) and risperidone response in Chinese schizophrenia patients. Wei Z, et al. Prog Neuropsychopharmacol Biol Psychiatry, 2009 Apr 30. PMID 19233240.