

Goat Anti-GPI / Neuroleukin (aa81-93) Antibody
Peptide-affinity purified goat antibody
Catalog # AF2221a**Specification****Goat Anti-GPI / Neuroleukin (aa81-93) Antibody - Product Information**

Application	WB
Primary Accession	P06744
Other Accession	NP_000166 , 2821
Reactivity	Human
Predicted	Pig
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	63147

Goat Anti-GPI / Neuroleukin (aa81-93) Antibody - Additional Information**Gene ID** 2821**Other Names**

Glucose-6-phosphate isomerase, GPI, 5.3.1.9, Autocrine motility factor, AMF, Neuroleukin, NLK, Phosphoglucose isomerase, PGI, Phosphohexose isomerase, PHI, Sperm antigen 36, SA-36, GPI

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-GPI / Neuroleukin (aa81-93) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-GPI / Neuroleukin (aa81-93) Antibody - Protein Information**Name** GPI {ECO:0000303|PubMed:2387591, ECO:0000312|HGNC:HGNC:4458}**Function**

In the cytoplasm, catalyzes the conversion of glucose-6-phosphate to fructose-6-phosphate, the second step in glycolysis, and the reverse reaction during gluconeogenesis (PubMed:28803808). Besides it's role as a glycolytic enzyme, also acts as a secreted cytokine: acts as an angiogenic factor (AMF) that stimulates endothelial cell motility (PubMed:<a

[11437381](http://www.uniprot.org/citations/11437381)). Acts as a neurotrophic factor, neuroleukin, for spinal and sensory neurons (PubMed:[3352745](http://www.uniprot.org/citations/3352745), PubMed:[11004567](http://www.uniprot.org/citations/11004567)). It is secreted by lectin-stimulated T-cells and induces immunoglobulin secretion (PubMed:[3352745](http://www.uniprot.org/citations/3352745), PubMed:[11004567](http://www.uniprot.org/citations/11004567)).

Cellular Location

Cytoplasm. Secreted

Goat Anti-GPI / Neuroleukin (aa81-93) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-GPI / Neuroleukin (aa81-93) Antibody - Images



AF2221a (0.3 µg/ml) staining of Human Heart lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



AF2221a (0.05 µg/ml) staining of HeLa (A), A431 (B), A549 (C), MCF7 (D), Jurkat (E) and K562 (F) lysates (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-GPI / Neuroleukin (aa81-93) Antibody - Background

This gene belongs to the GPI family whose members encode multifunctional phosphoglucose isomerase proteins involved in energy pathways. The protein encoded by this gene is a dimeric enzyme that catalyzes the reversible isomerization of glucose-6-phosphate and fructose-6-phosphate. The protein functions in different capacities inside and outside the cell. In the cytoplasm, the gene product is involved in glycolysis and gluconeogenesis, while outside the cell it functions as a neurotrophic factor for spinal and sensory neurons. Defects in this gene are the cause of nonspherocytic hemolytic anemia and a severe enzyme deficiency can be associated with hydrops fetalis, immediate neonatal death and neurological impairment.

Goat Anti-GPI / Neuroleukin (aa81-93) Antibody - References

New genetic associations detected in a host response study to hepatitis B vaccine. Davila S, et al. *Genes Immun*, 2010 Apr. PMID 20237496.

PI 3-kinase/Akt and STAT3 are required for the prevention of TGF-beta-induced Hep3B cell apoptosis by autocrine motility factor/phosphoglucose isomerase. Shih WL, et al. *Cancer Lett*, 2010 Apr 28. PMID 19819066.

Phosphoglucose isomerase/autocrine motility factor promotes melanoma cell migration through ERK activation dependent on autocrine production of interleukin-8. Araki K, et al. *J Biol Chem*, 2009 Nov 20. PMID 19801670.

Phosphoglucose isomerase enhances colorectal cancer metastasis. Tsutsumi S, et al. *Int J Oncol*, 2009 Nov. PMID 19787266.

Phosphoglucose isomerase/autocrine motility factor mediates epithelial and mesenchymal phenotype conversions in breast cancer. Funasaka T, et al. *Cancer Res*, 2009 Jul 1. PMID 19531650.