

SH2-B / SH2B1 (isoform2) Antibody (C-Term)
Peptide-affinity purified goat antibody
Catalog # AF2242a**Specification**

SH2-B / SH2B1 (isoform2) Antibody (C-Term) - Product Information

Application	E
Primary Accession	O9NRF2
Other Accession	NP_056318.2 , 25970 , 20399 (mouse)
Predicted	Human, Mouse, Rat, Pig, Dog, Cow
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	79366

SH2-B / SH2B1 (isoform2) Antibody (C-Term) - Additional Information**Gene ID** 25970**Other Names**

SH2B adapter protein 1, Pro-rich, PH and SH2 domain-containing signaling mediator, PSM, SH2 domain-containing protein 1B, SH2B1, KIAA1299, SH2B

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SH2-B / SH2B1 (isoform2) Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

SH2-B / SH2B1 (isoform2) Antibody (C-Term) - Protein Information**Name** SH2B1**Synonyms** KIAA1299, SH2B**Function**

Adapter protein for several members of the tyrosine kinase receptor family. Involved in multiple signaling pathways mediated by Janus kinase (JAK) and receptor tyrosine kinases, including the receptors of insulin (INS), insulin-like growth factor I (IGF1), nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), glial cell line-derived neurotrophic factor (GDNF), platelet-derived growth factor (PDGF) and fibroblast growth factors (FGFs). In growth hormone

(GH) signaling, autophosphorylated ('Tyr-813') JAK2 recruits SH2B1, which in turn is phosphorylated by JAK2 on tyrosine residues. These phosphotyrosines form potential binding sites for other signaling proteins. GH also promotes serine/threonine phosphorylation of SH2B1 and these phosphorylated residues may serve to recruit other proteins to the GHR-JAK2-SH2B1 complexes, such as RAC1. In leptin (LEP) signaling, binds to and potentiates the activation of JAK2 by globally enhancing downstream pathways. In response to leptin, binds simultaneously to both, JAK2 and IRS1 or IRS2, thus mediating formation of a complex of JAK2, SH2B1 and IRS1 or IRS2. Mediates tyrosine phosphorylation of IRS1 and IRS2, resulting in activation of the PI 3- kinase pathway. Acts as a positive regulator of NGF-mediated activation of the Akt/Forkhead pathway; prolongs NGF-induced phosphorylation of AKT1 on 'Ser-473' and AKT1 enzymatic activity. Enhances the kinase activity of the cytokine receptor-associated tyrosine kinase JAK2 and of other receptor tyrosine kinases, such as FGFR3 and NTRK1. For JAK2, the mechanism seems to involve dimerization of both, SH2B1 and JAK2. Enhances RET phosphorylation and kinase activity. Isoforms seem to be differentially involved in IGF-I and PDGF-induced mitogenesis (By similarity).

Cellular Location

Cytoplasm. Membrane. Nucleus. Note=Shuttles between the nucleus and the cytoplasm.

Tissue Location

Widely expressed with highest levels in skeletal muscle and ovary.

SH2-B / SH2B1 (isoform2) Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SH2-B / SH2B1 (isoform2) Antibody (C-Term) - Images**SH2-B / SH2B1 (isoform2) Antibody (C-Term) - Background**

This antibody is expected to recognise isoform 2 (NP_056318.2) only. Reported variants represent identical protein (NP_001139268.1; NP_056318.2; NP_001139284.1).

SH2-B / SH2B1 (isoform2) Antibody (C-Term) - References

SH2-B family members differentially regulate JAK family tyrosine kinases. O'Brien KB, O'Shea JJ, Carter-Su C. J Biol Chem 2002 Mar 8;277(10):8673-81 PMID: 11751854