

Neuro-d4 / DPF1 Antibody (C-Term)
Peptide-affinity purified goat antibody
Catalog # AF2313a**Specification**

Neuro-d4 / DPF1 Antibody (C-Term) - Product Information

Application	WB
Primary Accession	Q92782
Other Accession	NP_001128627.1 , NP_004638.2 , NP_001128628.1 , 8193 , 29861 (mouse) , 50545 (rat)
Reactivity	Human
Predicted	Mouse, Rat, Cow
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	44128

Neuro-d4 / DPF1 Antibody (C-Term) - Additional Information**Gene ID** 8193**Other Names**

Zinc finger protein neuro-d4, BRG1-associated factor 45B, BAF45B, D4, zinc and double PHD fingers family 1, DPF1, BAF45B, NEUD4

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Neuro-d4 / DPF1 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Neuro-d4 / DPF1 Antibody (C-Term) - Protein Information**Name** DPF1 ([HGNC:20225](#))**Synonyms** BAF45B, NEUD4**Function**

May have an important role in developing neurons by participating in regulation of cell survival, possibly as a neurospecific transcription factor. Belongs to the neuron-specific chromatin

remodeling complex (nBAF complex). During neural development a switch from a stem/progenitor to a postmitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to postmitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth (By similarity).

Cellular Location


Cytoplasm. Nucleus.

Neuro-d4 / DPF1 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Neuro-d4 / DPF1 Antibody (C-Term) - Images



AF2313a(1 µg/ml) staining of Human Brain (Cerebellum) lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Neuro-d4 / DPF1 Antibody (C-Term) - Background

This antibody is expected to recognize all three reported isoforms (NP_001128627.1; NP_004638.2; NP_001128628.1).

Neuro-d4 / DPF1 Antibody (C-Term) - References

The d4 gene family in the human genome. Chestkov AV, Baka ID, Kost MV, Georgiev GP, Buchman VL. Genomics. 1996 Aug 15;36(1):174-7. PMID: 8812431