

eotaxin Antibody (C-Term)
Peptide-affinity purified goat antibody
Catalog # AF2857a**Specification**

eotaxin Antibody (C-Term) - Product Information

Application	E
Primary Accession	P51671
Other Accession	NP_002977.1 , 6356
Predicted	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	10732

eotaxin Antibody (C-Term) - Additional Information**Gene ID** 6356**Other Names**

Eotaxin, C-C motif chemokine 11, Eosinophil chemotactic protein, Small-inducible cytokine A11, CCL11, SCYA11

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

eotaxin Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

eotaxin Antibody (C-Term) - Protein Information**Name** CCL11**Synonyms** SCYA11**Function**

In response to the presence of allergens, this protein directly promotes the accumulation of eosinophils, a prominent feature of allergic inflammatory reactions (PubMed:8597956). Binds to CCR3 (PubMed:8631813).

Cellular Location
Secreted.

eotaxin Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

eotaxin Antibody (C-Term) - Images

eotaxin Antibody (C-Term) - References

Differential regulation of CCL-11/eotaxin-1 and CXCL-8/IL-8 by gram-positive and gram-negative bacteria in human airway smooth muscle cells. Issa R, Sorrentino R, Sukkar MB, Sriskandan S, Chung KF, Mitchell JA. Respir Res. 2008 Apr 1;9:30. PMID: 18380907