

Sp1(mouse) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF2879a**Specification**

Sp1(mouse) Antibody (internal region) - Product Information

Application	WB
Primary Accession	O89090.2
Other Accession	NP_038700.2 , 20683 (mouse)
Reactivity	Mouse
Predicted	Rat, Pig
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

Sp1(mouse) Antibody (internal region) - Additional Information**Other Names**

Sp1; trans-acting transcription factor 1; 1110003E12Rik; AA450830; AI845540; Sp1-1

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Sp1(mouse) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Sp1(mouse) Antibody (internal region) - Protein Information**Sp1(mouse) Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Sp1(mouse) Antibody (internal region) - Images



AF2879a (0.3 µg/ml) staining of NIH3T3 nuclear lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Sp1(mouse) Antibody (internal region) - References

Large-scale mutagenesis in p19(ARF)- and p53-deficient mice identifies cancer genes and their collaborative networks. Uren AG, Kool J, Matentzoglou K, de Ridder J, Mattison J, van Uitert M, Lagcher W, Sie D, Tanger E, Cox T, Reinders M, Hubbard TJ, Rogers J, Jonkers J, Wessels L, Adams DJ, van Lohuizen M, Berns A, Cell 2008 May 133 (4): 727-41. PMID: 18485879