

CDC2L6 Antibody (C-Term)
Peptide-affinity purified goat antibody
Catalog # AF3371a**Specification**

CDC2L6 Antibody (C-Term) - Product Information

Application	IF, FC
Primary Accession	O9BWU1
Other Accession	NP_055891.1 , 23097 , 78334 (mouse) , 309804 (rat)
Predicted Host	Human, Mouse, Rat, Dog, Cow
Clonality	Goat
Concentration	Polyclonal
Isotype	0.5 mg/ml
Calculated MW	IgG
	56802

CDC2L6 Antibody (C-Term) - Additional Information**Gene ID** 23097**Other Names**

Cyclin-dependent kinase 19, 2.7.11.22, CDC2-related protein kinase 6, Cell division cycle 2-like protein kinase 6, Cell division protein kinase 19, Cyclin-dependent kinase 11, Death-preventing kinase, CDK19, CDC2L6, CDK11, KIAA1028

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CDC2L6 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

CDC2L6 Antibody (C-Term) - Protein Information**Name** CDK19**Synonyms** CDC2L6, CDK11, KIAA1028**Cellular Location**


Cytoplasm. Cytoplasm, perinuclear region. Nucleus

CDC2L6 Antibody (C-Term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CDC2L6 Antibody (C-Term) - Images



EB10476 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing vesicle and weak nuclear staining. The nuclear sta



EB10476 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) fol

CDC2L6 Antibody (C-Term) - References

Cyclin-dependent kinases: a family portrait. Malumbres M, Harlow E, Hunt T, Hunter T, Lahti JM, Manning G, Morgan DO, Tsai LH, Wolgemuth DJ. Nat Cell Biol. 2009 Nov;11(11):1275-6. PMID: 19884882