

Mucin 19 / Apomucin Antibody (Internal region, near N-Term)
Peptide-affinity purified goat antibody
Catalog # AF3429a

Specification

Mucin 19 / Apomucin Antibody (Internal region, near N-Term) - Product Information

Application	E
Primary Accession	Q6PZE0.2
Other Accession	NP_997126.2 , 239611 (mouse) , 497227 (rat)
Predicted	Mouse, Rat
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

Mucin 19 / Apomucin Antibody (Internal region, near N-Term) - Additional Information

Other Names

Muc19; mucin 19; MUC-19; gel-forming secreted mucin-19; mucin-19; sublingual apomucin; mucin apoprotein

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mucin 19 / Apomucin Antibody (Internal region, near N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mucin 19 / Apomucin Antibody (Internal region, near N-Term) - Protein Information

Mucin 19 / Apomucin Antibody (Internal region, near N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mucin 19 / Apomucin Antibody (Internal region, near N-Term) - Images**Mucin 19 / Apomucin Antibody (Internal region, near N-Term) - References**

Detailed haplotype-tagging study of germline variation of MUC19 in inflammatory bowel disease. Phillips AM, Nimmo ER, Van Limbergen J, Drummond HE, Smith L, Satsangi J, Inflammatory bowel diseases 2010 Apr 16 (4): 557-8. PMID: 19714762