

Muc19 / Smgc Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3434a**Specification**

Muc19 / Smgc Antibody (internal region) - Product Information

Application	ICC
Primary Accession	O6JHY2.1
Other Accession	NP_945121.1 , 223809 (mouse)
Reactivity	Mouse
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

Muc19 / Smgc Antibody (internal region) - Additional Information**Other Names**

Smgc; submandibular gland protein C; 2310010P21Rik; DXImx49e; Muc19; Sfc21; OTTMUSP00000022832; neonatal submandibular gland protein C

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.


Precautions

Muc19 / Smgc Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Muc19 / Smgc Antibody (internal region) - Protein Information**Muc19 / Smgc Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Muc19 / Smgc Antibody (internal region) - Images

AF3434a (4ug/ml) staining of Mouse SubMandibular Gland cells at E18. Nuclear counterstain with DAPI in green.

Muc19 / Smgc Antibody (internal region) - References

Expression of Muc19/Smgc gene products during murine sublingual gland development: cytodifferentiation and maturation of salivary mucous cells. Das B, Cash MN, Hand AR, Shivazad A, Culp DJ, The journal of histochemistry and cytochemistry : official journal of the Histochemistry Society 2009 Apr 57 (4): 383-96. PMID: 19110483