

Npc2 (mouse) (aa21-33) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3496a**Specification**

Npc2 (mouse) (aa21-33) Antibody (internal region) - Product Information

Application	WB
Primary Accession	O9Z0J0.1
Other Accession	NP_075898.1 , 67963 (mouse) , 286898 (rat)
Reactivity	Mouse
Predicted	Rat
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

Npc2 (mouse) (aa21-33) Antibody (internal region) - Additional Information**Other Names**

Npc2; Niemann-Pick disease, type C2; 2700012J19Rik; AA408070; AU045843; HE1; epididymal secretory protein E1; mE1; niemann Pick type C2 protein homolog

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Npc2 (mouse) (aa21-33) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.


Npc2 (mouse) (aa21-33) Antibody (internal region) - Protein Information**Npc2 (mouse) (aa21-33) Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

Npc2 (mouse) (aa21-33) Antibody (internal region) - Images



AF3496a (0.1 µg/ml) staining of Mouse Kidney lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Npc2 (mouse) (aa21-33) Antibody (internal region) - References

Somatic cell plasticity and Niemann-pick type C2 protein: adipocyte differentiation and function.
Csepegi C, Jiang M, Frolov A, The Journal of biological chemistry 2010 Sep 285 (39): 30347-54.
PMID: 20650896