

KDM2A (aa445-459) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3534a**Specification**

KDM2A (aa445-459) Antibody (internal region) - Product Information

Application	WB
Primary Accession	O9Y2K7
Other Accession	NP_036440.1 , 22992
Reactivity	Human
Predicted	Mouse, Rat, Pig, Dog, Cow
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	132793

KDM2A (aa445-459) Antibody (internal region) - Additional Information**Gene ID** 22992**Other Names**

Lysine-specific demethylase 2A, 1.14.11.27, CXXC-type zinc finger protein 8, F-box and leucine-rich repeat protein 11, F-box protein FBL7, F-box protein Lilina, F-box/LRR-repeat protein 11, JmjC domain-containing histone demethylation protein 1A, [Histone-H3]-lysine-36 demethylase 1A, KDM2A, CXXC8, FBL7, FBXL11, JHDM1A, KIAA1004

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

KDM2A (aa445-459) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

KDM2A (aa445-459) Antibody (internal region) - Protein Information**Name** KDM2A**Function**

Histone demethylase that specifically demethylates 'Lys-36' of histone H3, thereby playing a central role in histone code. Preferentially demethylates dimethylated H3 'Lys-36' residue while it has weak or no activity for mono- and tri-methylated H3 'Lys-36'. May also recognize and bind to some phosphorylated proteins and promote their ubiquitination and degradation. Required to

maintain the heterochromatic state. Associates with centromeres and represses transcription of small non-coding RNAs that are encoded by the clusters of satellite repeats at the centromere. Required to sustain centromeric integrity and genomic stability, particularly during mitosis. Regulates circadian gene expression by repressing the transcriptional activator activity of CLOCK-BMAL1 heterodimer and RORA in a catalytically- independent manner (PubMed:26037310).

Cellular Location

Nucleus, nucleoplasm. Chromosome Note=Punctate expression throughout the nucleoplasm and enriched in the perinucleolar region (PubMed:19001877, PubMed:20417597). Specifically nucleates at CpG islands where it's presence results in chromatin depleted in H3K36me2 (PubMed:19001877, PubMed:20417597)

Tissue Location

Widely expressed, with highest levels in brain, testis and ovary, followed by lung.

KDM2A (aa445-459) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KDM2A (aa445-459) Antibody (internal region) - Images



AF3534a (0.5 µg/ml) staining of Human Uterus lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

KDM2A (aa445-459) Antibody (internal region) - References

Prediction of the coding sequences of unidentified human genes. XIII. The complete sequences of

100 new cDNA clones from brain which code for large proteins in vitro. Nagase T, Ishikawa K, Suyama M, Kikuno R, Hirosawa M, Miyajima N, Tanaka A, Kotani H, Nomura N, Ohara O. DNA Res. 1999 Feb 26;6(1):63-70. PMID: 10231032