

Dbf4 (mouse) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3545a**Specification**

Dbf4 (mouse) Antibody (internal region) - Product Information

Application	WB
Primary Accession	O9OZ41.1
Other Accession	NP_038754.1 , NP_001177646.1 , 27214 (mouse) , 312046 (rat)
Reactivity	Mouse, Rat
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

Dbf4 (mouse) Antibody (internal region) - Additional Information**Other Names**

Dbf4; DBF4 homolog (S. cerevisiae); AA545217; Ask; activator of S phase kinase; protein DBF4 homolog A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Dbf4 (mouse) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.


Dbf4 (mouse) Antibody (internal region) - Protein Information**Dbf4 (mouse) Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

Dbf4 (mouse) Antibody (internal region) - Images



AF3545a (1 µg/ml) staining of Mouse Testis lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Dbf4 (mouse) Antibody (internal region) - Background

This antibody is expected to recognize both reported isoforms (NP_038754.1; NP_001177646.1).

Dbf4 (mouse) Antibody (internal region) - References

The role of Dbf4/Drf1-dependent kinase Cdc7 in DNA-damage checkpoint control. Tsuji T, Lau E, Chiang GG, Jiang W. Mol Cell. 2008 Dec 26;32(6):862-9. PMID: 19111665