

Wfdc1 (mouse) Antibody (internal region , near N-Term)
Peptide-affinity purified goat antibody
Catalog # AF3588a**Specification**

Wfdc1 (mouse) Antibody (internal region , near N-Term) - Product Information

Application	WB
Primary Accession	Q9ESH5.1
Other Accession	NP_075884.1 , 67866 (mouse) , 171112 (rat)
Reactivity	Mouse
Predicted	Rat
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

Wfdc1 (mouse) Antibody (internal region , near N-Term) - Additional Information**Other Names**

WFDC1; WAP four-disulfide core domain 1; ps20; WAP four-disulfide core domain 1 homolog; WAP four-disulfide core domain protein 1; prostate stromal protein ps20; ps20 growth inhibitor; 2310058A03Rik

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Wfdc1 (mouse) Antibody (internal region , near N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Wfdc1 (mouse) Antibody (internal region , near N-Term) - Protein Information**Wfdc1 (mouse) Antibody (internal region , near N-Term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

Wfdc1 (mouse) Antibody (internal region , near N-Term) - Images

AF3588a (0.3 µg/ml) staining of Mouse Heart lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Wfdc1 (mouse) Antibody (internal region , near N-Term) - References

Molecular analysis of WFDC1/ps20 gene in prostate cancer. Watson JE, Kamkar S, James K, Kowbel D, Andaya A, Paris PL, Simko J, Carroll P, McAlhany S, Rowley D, Collins C. Prostate. 2004 Oct 1;61(2):192-9. PMID: 15305342