10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Terminal Uridylyltransferase 4 (mouse) Antibody (internal region)

Peptide-affinity purified goat antibody
Catalog # AF3614a

Specification

Terminal Uridylyltransferase 4 (mouse) Antibody (internal region) - Product Information

Application	**WB**
Primary Accession	B2RX14.2
Other Accession	NP_780681.2, 230594 (mouse), 313481 (rat)
Reactivity	**Mouse, Rat**
Host	**Goat**
Clonality	**Polyclonal**
Concentration	**0.5 mg/ml**
Isotype	**IgG**

Terminal Uridylyltransferase 4 (mouse) Antibody (internal region) - Additional Information

Other Names
Zcchc11; zinc finger, CCHC domain containing 11; RP23-406B13.1; 6030404K05Rik; 9230115F04Rik; KIAA0191; PPAPD3; mKIAA0191; OTTMUSP00000008601; TUTase 4; terminal uridylyltransferase 4; zinc finger CCHC domain-containing protein 11

Format
0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
Terminal Uridylyltransferase 4 (mouse) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Terminal Uridylyltransferase 4 (mouse) Antibody (internal region) - Protein Information

Terminal Uridylyltransferase 4 (mouse) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

- Flow Cytomety
- Cell Culture

Terminal Uridylyltransferase 4 (mouse) Antibody (internal region) - Images



AF3614a (0.05 µg/ml) staining of Rat Brain lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



AF3614a (0.2 µg/ml) staining of different Testis lysates from wildtype 8-9 week old C57BL/6 (wt) and knock-out (KO) mice (35 µg protein in RIPA buffer). Primary incubation was 2 hours. Detected by chemiluminescence. Data obtained from an anonymous customer.

Terminal Uridylyltransferase 4 (mouse) Antibody (internal region) - References

A novel Zinc finger protein, ZCCHC11, interacts with TIFA and modulates TLR signaling. Minoda Y, Saeki K, Aki D, Takaki H, Sanada T, Koga K, Kobayashi T, Takaesu G, Yoshimura A. Biochem Biophys Res Commun. 2006 Jun 9;344(3):1023-30. PMID: 16643855