

Muc16 (Rat) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3621a**Specification**

Muc16 (Rat) Antibody (internal region) - Product Information

Application	E
Other Accession	XP_235886.5 , 73732 (mouse) , 315451 (rat)
Predicted	Mouse, Rat
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

Muc16 (Rat) Antibody (internal region) - Additional Information**Other Names**

MUC16; mucin 16, cell surface associated; CA125; CA-125; CA125 ovarian cancer antigen; MUC-16; mucin-16; ovarian cancer-related tumor marker CA125; ovarian carcinoma antigen CA125; RGD1308422; 1110008I14Rik; Gm1480

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Muc16 (Rat) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Muc16 (Rat) Antibody (internal region) - Protein Information**Muc16 (Rat) Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Muc16 (Rat) Antibody (internal region) - Images**Muc16 (Rat) Antibody (internal region) - References**

Serum HE4 concentration differentiates malignant ovarian tumours from ovarian endometriotic cysts. Huhtinen K, Suvitie P, Hiissa J, Junnila J, Huvila J, Kujari H, Setälä M, Härkki P, Jalkanen J, Fraser J, Mäkinen J, Auranen A, Poutanen M, Perheentupa A. Br J Cancer. 2009 Apr 21;100(8):1315-9. PMID: 19337252