10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CDH1 (aa662-675) Antibody (internal region)

Peptide-affinity purified goat antibody
Catalog # AF3666a

Specification

CDH1 (aa662-675) Antibody (internal region) - Product Information

Application	**WB**
Primary Accession	P12830
Other Accession	NP_004351.1, 999, 83502 (rat)
Reactivity	**Human, Mouse, Rat, Pig**
Predicted	**Dog**
Host	**Goat**
Clonality	**Polyclonal**
Concentration	**0.5 mg/ml**
Isotype	**IgG**
Calculated MW	**97456**

CDH1 (aa662-675) Antibody (internal region) - Additional Information

Gene ID 999

Other Names
Cadherin-1, CAM 120/80, Epithelial cadherin, E-cadherin, Uvomorulin, CD324, E-Cad/CTF1, E-Cad/CTF2, E-Cad/CTF3, CDH1, CDHE, UVO

Format
0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
CDH1 (aa662-675) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

CDH1 (aa662-675) Antibody (internal region) - Protein Information

Name CDH1

Synonyms CDHE, UVO

Function
Cadherins are calcium-dependent cell adhesion proteins (PubMed:11976333). They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. CDH1 is involved in mechanisms

regulating cell-cell adhesions, mobility and proliferation of epithelial cells (PubMed:11976333). Has a potent invasive suppressor role. It is a ligand for integrin alpha-E/beta-7.

Cellular Location
Cell junction, adherens junction. Cell membrane; Single-pass type I membrane protein. Endosome. Golgi apparatus, trans-Golgi network. Note=Colocalizes with DLGAP5 at sites of cell-cell contact in intestinal epithelial cells. Anchored to actin microfilaments through association with alpha-, beta- and gamma-catenin. Sequential proteolysis induced by apoptosis or calcium influx, results in translocation from sites of cell-cell contact to the cytoplasm Colocalizes with RAB11A endosomes during its transport from the Golgi apparatus to the plasma membrane


Tissue Location
Non-neural epithelial tissues.

CDH1 (aa662-675) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CDH1 (aa662-675) Antibody (internal region) - Images



AF3666a (0.5 µg/ml) staining of Human Kidney lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

CDH1 (aa662-675) Antibody (internal region) - Background

The immunizing peptide represents part of the extracellular domain.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CDH1 (aa662-675) Antibody (internal region) - References

E-cadherin expression on human carcinoma cell affects trastuzumab-mediated antibody-dependent cellular cytotoxicity through killer cell lectin-like receptor G1 on natural killer cells. Yamauchi C, Fujii S, Kimura T, Kuwata T, Wada N, Mukai H, Matsumoto N, Fukayama M, Ochiai A. Int J Cancer. 2011 May 1;128(9):2125-37 PMID: 21387286