

Hsp31p (yeast, aa50-64) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3773a**Specification**

Hsp31p (yeast, aa50-64) Antibody (internal region) - Product Information

Application	WB
Primary Accession	Q04432.1
Other Accession	NP_010822.1
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

Hsp31p (yeast, aa50-64) Antibody (internal region) - Additional Information**Other Names**

Hsp31p; YDR533C

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions


Hsp31p (yeast, aa50-64) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Hsp31p (yeast, aa50-64) Antibody (internal region) - Protein Information**Hsp31p (yeast, aa50-64) Antibody (internal region) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Hsp31p (yeast, aa50-64) Antibody (internal region) - Images



AF3773a (0.05 µg/ml) staining of yeast lysate (wt) and the KO (delta) in the left panel and the Ponceau stain in the right panel (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence. Data obtained by Dr. M. Skoneczny, IBB, Warsaw, Poland.



EB11539 (0.05 µg/ml) staining of yeast lysate (wt) and the KO (delta) in the left panel and the Ponceau stain in the right panel (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence. Data obtained by Dr. M. Skoneczny

Hsp31p (yeast, aa50-64) Antibody (internal region) - References

: . PMID: