

DNAJC5G (human) Antibody (internal region, near C-Term)
Peptide-affinity purified goat antibody
Catalog # AF3895a

Specification

DNAJC5G (human) Antibody (internal region, near C-Term) - Product Information

Application	IHC
Primary Accession	Q8N7S2
Other Accession	NP_775921.1 , 285126
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	21433

DNAJC5G (human) Antibody (internal region, near C-Term) - Additional Information

Gene ID 285126

Other Names

DnaJ homolog subfamily C member 5G, Cysteine string protein-gamma, CSP-gamma, DNAJC5G

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DNAJC5G (human) Antibody (internal region, near C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

DNAJC5G (human) Antibody (internal region, near C-Term) - Protein Information

Name DNAJC5G

Cellular Location

Membrane; Lipid-anchor

Tissue Location


Testis specific.

DNAJC5G (human) Antibody (internal region, near C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DNAJC5G (human) Antibody (internal region, near C-Term) - Images



AF3895a (5 µg/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

DNAJC5G (human) Antibody (internal region, near C-Term) - References

: . PMID: