

BEND5 Antibody (C-Term)
Peptide-affinity purified goat antibody
Catalog # AF3996a**Specification**

BEND5 Antibody (C-Term) - Product Information

Application	WB
Primary Accession	Q7L4P6
Other Accession	NP_078879.2 , 79656 , 67621 (mouse) , 362564 (rat)
Reactivity	Human, Mouse
Predicted	Rat, Dog, Cow
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	48182

BEND5 Antibody (C-Term) - Additional Information**Gene ID** 79656**Other Names**

BEN domain-containing protein 5, BEND5, C1orf165

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BEND5 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

BEND5 Antibody (C-Term) - Protein Information**Name** BEND5**Synonyms** C1orf165**Function**


Acts as a transcriptional repressor (PubMed:23468431).

BEND5 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BEND5 Antibody (C-Term) - Images



AF3996a (0.3 µg/ml) staining of Daudi (A), Human Tonsil (B), Human Spleen (C) and Mouse Spleen (D) lysates (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

BEND5 Antibody (C-Term) - References

Transcriptional programs in transient embryonic zones of the cerebral cortex defined by high-resolution mRNA sequencing. Ayoub AE, Oh S, Xie Y, Leng J, Cotney J, Dominguez MH, Noonan JP, Rakic P. Proceedings of the National Academy of Sciences of the United States of America 2011 Sep 108 (36): 14950-5. PMID: 21873192