

contactin 1 (aa585-870) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF4042a

Specification

contactin 1 (aa585-870) Antibody (internal region) - Product Information

Application	WB, IHC
Primary Accession	Q12860
Other Accession	NP_001834.2 , NP_778203.1 , 1272 , 12805 (mouse) , 117258 (rat)
Reactivity	Human, Mouse, Rat
Predicted	Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	113320

contactin 1 (aa585-870) Antibody (internal region) - Additional Information

Gene ID 1272

Other Names

Contactin-1, Glycoprotein gp135, Neural cell surface protein F3, CNTN1

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

contactin 1 (aa585-870) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

contactin 1 (aa585-870) Antibody (internal region) - Protein Information

Name CNTN1

Function

Contactins mediate cell surface interactions during nervous system development. Involved in the formation of paranodal axo-glial junctions in myelinated peripheral nerves and in the signaling between axons and myelinating glial cells via its association with CNTNAP1. Participates in oligodendrocytes generation by acting as a ligand of NOTCH1. Its association with NOTCH1 promotes NOTCH1 activation through the released notch intracellular domain (NICD) and subsequent translocation to the nucleus. Interaction with TNR induces a repulsion of neurons and

an inhibition of neurite outgrowth (By similarity).

Cellular Location

[Isoform 1]: Cell membrane; Lipid-anchor, GPI- anchor; Extracellular side


Tissue Location

Strongly expressed in brain and in neuroblastoma and retinoblastoma cell lines. Lower levels of expression in lung, pancreas, kidney and skeletal muscle.


contactin 1 (aa585-870) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

contactin 1 (aa585-870) Antibody (internal region) - Images

AF4042a (1 µg/ml) staining of Human Cerebellum lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



AF4042a (5 µg/ml) staining of paraffin embedded Human Cerebellum. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

contactin 1 (aa585-870) Antibody (internal region) - Background

This antibody is expected to recognize isoform 1 (NP_001834.2) and isoform 2 (NP_778203.1).

contactin 1 (aa585-870) Antibody (internal region) - References

A complex between contactin-1 and the protein tyrosine phosphatase PTPRZ controls the development of oligodendrocyte precursor cells. Lamprianou S, Chatzopoulou E, Thomas JL, Bouyain S, Harroch S. Proc Natl Acad Sci U S A. 2011 Oct 18;108(42):17498-503. PMID: 21969550