

**Goat Anti-DC-SIGN / CD209 Antibody (internal region)**  
**Purified Goat Polyclonal Antibody**  
**Catalog # AF4236a****Specification**

---

**Goat Anti-DC-SIGN / CD209 Antibody (internal region) - Product Information**

Application	WB
Primary Accession	<a href="#">O9NNX6</a>
Other Accession	<a href="#">NP_066978.1</a> , <a href="#">NP_001138368.1</a> , <a href="#">NP_001138369.1</a> , <a href="#">NP_001138365.1</a> , <a href="#">NP_001138366.1</a> , <a href="#">NP_001138367.1</a> , <a href="#">NP_001138371.1</a>
Reactivity	Human
Predicted	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5
Calculated MW	45775

**Goat Anti-DC-SIGN / CD209 Antibody (internal region) - Additional Information****Gene ID** 30835**Other Names**

CD209; CD209 molecule; CDSIGN; CLEC4L; DC-SIGN; DC-SIGN1; C-type lectin domain family 4 member L; C-type lectin domain family 4, member L; CD209 antigen; HIV gp120-binding protein; dendritic cell-specific ICAM-3-grabbing non-integrin 1; dendritic cell-spe

**Format**

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

**Immunogen**

Peptide with sequence C-NRGEPNNVGEED, from the internal region of the protein sequence according to [NP\\_066978.1](#); [NP\\_001138368.1](#); [NP\\_001138369.1](#); [NP\\_001138365.1](#); [NP\\_001138366.1](#); [NP\\_001138367.1](#); [NP\\_001138371.1](#).

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-DC-SIGN / CD209 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-DC-SIGN / CD209 Antibody (internal region) - Protein Information**

**Name** CD209

**Synonyms** CLEC4L

**Function**

Pathogen-recognition receptor expressed on the surface of immature dendritic cells (DCs) and involved in initiation of primary immune response. Thought to mediate the endocytosis of pathogens which are subsequently degraded in lysosomal compartments. The receptor returns to the cell membrane surface and the pathogen-derived antigens are presented to resting T-cells via MHC class II proteins to initiate the adaptive immune response.

**Cellular Location**

[Isoform 1]: Cell membrane; Single- pass type II membrane protein [Isoform 3]: Cell membrane; Single- pass type II membrane protein [Isoform 5]: Cell membrane; Single- pass type II membrane protein [Isoform 7]: Secreted. [Isoform 9]: Secreted. [Isoform 11]: Secreted.

**Tissue Location**

Predominantly expressed in dendritic cells and in DC-residing tissues. Also found in placental macrophages, endothelial cells of placental vascular channels, peripheral blood mononuclear cells, and THP-1 monocytes.

**Goat Anti-DC-SIGN / CD209 Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Goat Anti-DC-SIGN / CD209 Antibody (internal region) - Images**



AF4236a (0.3 µg/ml) staining of Human Bone Marrow lysate (35 µg protein in RIPA buffer).

Primary incubation was 1 hour. Detected by chemiluminescence.

**Goat Anti-DC-SIGN / CD209 Antibody (internal region) - References**

Role of DC-SIGN in Lassa virus entry into human dendritic cells. Goncalves AR, Moraz ML, Pasquato A, Helenius A, Lozach PY, Kunz S. Journal of virology 2013 Nov 87 (21): 11504-15.