10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CD1a (Mature Langerhans Cells Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone SPM120]

Catalog # AH10832

Specification

CD1a (Mature Langerhans Cells Marker) Antibody - With BSA and Azide - Product Information

Application	**,14,3,4,**
Primary Accession	P06126
Other Accession	909, 1309
Reactivity	**Human**
Host	**Mouse**
Clonality	**Monoclonal**
Isotype	**Mouse / IgG1, kappa**
Calculated MW	**49kDa KDa**

CD1a (Mature Langerhans Cells Marker) Antibody - With BSA and Azide - Additional Information

Gene ID 909

Other Names
T-cell surface glycoprotein CD1a, T-cell surface antigen T6/Leu-6, hTa1 thymocyte antigen, CD1a, CD1A

Format
200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage
Store at 2 to 8°C.Antibody is stable for 24 months.

Precautions
CD1a (Mature Langerhans Cells Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

CD1a (Mature Langerhans Cells Marker) Antibody - With BSA and Azide - Protein Information

Name CD1A

Function
Antigen-presenting protein that binds self and non-self lipid and glycolipid antigens and presents them to T-cell receptors on natural killer T-cells.

Cellular Location
Cell membrane; Single-pass type I membrane protein. Membrane raft; Single-pass type I membrane protein. Endosome membrane; Single- pass type I membrane protein. Note=Subject to

intracellular trafficking between the cell membrane and endosomes (PubMed:11231314). Localizes to cell surface lipid rafts (PubMed:18178838).


Tissue Location
Expressed on cortical thymocytes, epidermal Langerhans cells, dendritic cells, on certain T-cell leukemias, and in various other tissues.

CD1a (Mature Langerhans Cells Marker) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CD1a (Mature Langerhans Cells Marker) Antibody - With BSA and Azide - Images



Formalin-fixed, paraffin-embedded human Skin stained with CD1a Monoclonal Antibody (SPM120).

CD1a (Mature Langerhans Cells Marker) Antibody - With BSA and Azide - Background

At least five CD1 genes (CD1a, b, c, d, and e) are identified. CD1 proteins have been demonstrated to restrict T cell response to non-peptide lipid and glycolipid antigens and play a role in non-classical antigen presentation. CD1a is a non-polymorphic MHC Class 1 related cell surface glycoprotein, expressed in association with Beta-2 microglobulin. Anti-CD1a labels Langerhans cell histiocytosis (Histiocytosis X), extranodal histiocytic sarcoma, a subset of T-lymphoblastic lymphoma/leukemia, and interdigitating dendritic cell sarcoma of the lymph node. ĀWhen combined with antibodies against TTF-1 and CD5, anti-CD1a is useful in distinguishing between pulmonary and thymic neoplasms since CD1a is consistently expressed in thymic lymphocytes in both typical and atypical thymomas, but only focally in 1/6 of thymic carcinomas and not in lymphocytes in pulmonary neoplasms. Anti-CD1a is reported to be a new marker for perivascular epithelial cell tumor (PEComa).Ā

CD1a (Mature Langerhans Cells Marker) Antibody - With BSA and Azide - References

McNally, AK. et al. 2011. Exp. Mol. Pathol. 91: 673-681. | Matsuda, A. et al. 2009. Invest. Ophthalmol. Vis. Sci. 50: 2871-2877. | Gulubova, M. et al. 2008. Clin. Exp. Metastasis. 25: 777-785. | Cassaday, RD. et al. 2007. Clin. Cancer Res. 13: 540-549.