

Phosphotyrosine (P-Tyr) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone SPM102]
Catalog # AH10927

Specification

Phosphotyrosine (P-Tyr) Antibody - With BSA and Azide - Product Information

Application ,1,14,3,4,5,
Host Mouse
Clonality Monoclonal
Isotype Mouse / IgG2b

Calculated MW Depends upon the phosphorylated target

KDa

Phosphotyrosine (P-Tyr) Antibody - With BSA and Azide - Additional Information

Format

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

Phosphotyrosine (P-Tyr) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

Phosphotyrosine (P-Tyr) Antibody - With BSA and Azide - Protein Information


Phosphotyrosine (P-Tyr) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Phosphotyrosine (P-Tyr) Antibody - With BSA and Azide - Images





Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with Phosphotyrosine Monoclonal Antibody (SPM102).

Phosphotyrosine (P-Tyr) Antibody - With BSA and Azide - Background

Protein phosphorylation is a fundamental event in the regulation of a large number of intracellular processes. Phosphorylation of specific tyrosine residues is the result of activation or stimulation of their respective protein tyrosine kinases. The phosphorylated proteins can be auto-phosphorylated kinases or certain cellular protein substrates. Tyrosine-phosphorylated proteins are involved in signal transduction and in the regulation of cell proliferation. Antibody to phosphotyrosine provides an excellent tool for the detection, characterization, and purification of phosphotyrosine containing proteins. This MAb shows no cross-reaction with other phosphoamino acids and is superb for multiple applications including staining of formalin/paraffin tissues.

Phosphotyrosine (P-Tyr) Antibody - With BSA and Azide - References

Glenny, Jr. JR, et. al. Journal of Immunological Methods, 1988, 109:277-85