

Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide Mouse Monoclonal Antibody [Clone CDH16/1071] Catalog # AH10983

Specification

Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide - Product Information

Application ,2,3,4,
Primary Accession 075309
Other Accession 1014, 513660

Reactivity Human, Mouse, Rat, Rabbit, Dog

Host Mouse Clonality Monoclonal

Isotype Mouse / IgG1, kappa

Calculated MW 130kDa KDa

Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide - Additional Information

Gene ID 1014

Other Names

Cadherin-16, Kidney-specific cadherin, Ksp-cadherin, CDH16

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide - Protein Information

Name CDH16


Function

Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types.

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location Kidney specific.






Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide - Images



Formalin-fixed, paraffin-embedded human Renal Cell Carcinoma stained with KSP-Cadherin Monoclonal Antibody (CDH16/1071)



Formalin-fixed, paraffin-embedded Rat Kidney stained with KSP-Cadherin Monoclonal Antibody (CDH16/1071)

Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide - Background

This MAb recognizes a protein of 130kDa, identified as Ksp-cadherin. Cadherins form a superfamily of related glycoproteins that mediate calcium-dependent cell adhesion and transmit signals from the extracellular matrix to the cytoplasm. Cadherins have been implicated in embryogenesis, tissue morphogenesis, tissue structure maintenance, cell polarization, neoplastic invasiveness and metastasis, and membrane transport. It is suggested that Ksp-cadherin is a marker for terminal





Tel: 858.875.1900 Fax: 858.875.1999

differentiation of the basolateral membranes of renal tubular epithelial cells. Within the kidney, Ksp-Cadherin is found exclusively in the basolateral membrane of renal tubular epithelial cells and collecting duct cells, and not in glomeruli, renal interstitial cells, or blood vessels. ĀKsp-Cadherin has been suggested to distinguish Chromophobe Renal-Cell Carcinoma from Oncocytoma.

Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide -References

Shen, S.S., et al. 2005. Kidney-specific cadherin, a specific marker for the distal portion of the nephron and related renal neoplasms. Mod Pathol. 18: 933-40. | Mazal, P.R., et al. 2005. Expression of kidney-specific cadherin distinguishes chromophobe renal cell carcinoma from renal oncocytoma. Hum Pathol. 36: 22-28. |