

---

**HCG-beta (Pregnancy & Choriocarcinoma Marker) Antibody - With BSA and Azide**  
**Mouse Monoclonal Antibody [Clone HCGb/211 ]**  
**Catalog # AH11069**

**Specification**

---

**HCG-beta (Pregnancy & Choriocarcinoma Marker) Antibody - With BSA and Azide -  
Product Information**

Application	''
Primary Accession	<a href="#">P01233</a>
Other Accession	<a href="#">1082</a> , <a href="#">172944</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1
Calculated MW	22kDa kDa

**HCG-beta (Pregnancy & Choriocarcinoma Marker) Antibody - With BSA and Azide -  
Additional Information**

**Other Names**

Choriogonadotropin subunit beta, CG-beta, Chorionic gonadotrophin chain beta, CGB, CGB3

**Storage**

Store at 2 to 8°C. Antibody is stable for 24 months.

**Precautions**

HCG-beta (Pregnancy & Choriocarcinoma Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

**HCG-beta (Pregnancy & Choriocarcinoma Marker) Antibody - With BSA and Azide -  
Protein Information**

**HCG-beta (Pregnancy & Choriocarcinoma Marker) Antibody - With BSA and Azide -  
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**HCG-beta (Pregnancy & Choriocarcinoma Marker) Antibody - With BSA and Azide -**

## Images

### **HCG-beta (Pregnancy & Choriocarcinoma Marker) Antibody - With BSA and Azide - Background**

This MAb reacts with a protein of 22kDa, identified as  $\beta$  sub-unit of HCG. It does not cross react with the  $\alpha$  sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the  $\alpha$  and  $\beta$  subunits. The  $\beta$  subunit is identical to that of thyroid stimulating hormone (TSH), follicle stimulating hormone (FSH), and luteinizing hormone (LH). hCG MAb detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. hCG expression by non-trophoblastic tumors may indicate aggressive behavior.

### **HCG-beta (Pregnancy & Choriocarcinoma Marker) Antibody - With BSA and Azide - References**

Cocquebert M et. al. Am J Physiol Endocrinol Metab. 2012;303(8):E950-8