10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Glycophorin A / CD235a (Erythrocyte Marker) Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone SPM599]
Catalog # AH11382

Specification

Glycophorin A / CD235a (Erythrocyte Marker) Antibody - With BSA and Azide - Product Information

Application	**,2,3,4,**
Primary Accession	P02724
Other Accession	2993, 2994, 434973, 654368
Reactivity	**Human**
Host	**Mouse**
Clonality	**Monoclonal**
Isotype	**Mouse / IgG1, kappa**
Calculated MW	**39kDa KDa**

Glycophorin A / CD235a (Erythrocyte Marker) Antibody - With BSA and Azide - Additional Information

Gene ID 2993

Other Names
Glycophorin-A, MN sialoglycoprotein, PAS-2, Sialoglycoprotein alpha, CD235a, GYPA, GPA

Storage
Store at 2 to 8°C.Antibody is stable for 24 months.

Precautions
Glycophorin A / CD235a (Erythrocyte Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

Glycophorin A / CD235a (Erythrocyte Marker) Antibody - With BSA and Azide - Protein Information

Name GYPA (HGNC:4702)

Synonyms GPA

Function
Component of the ankyrin-1 complex, a multiprotein complex involved in the stability and shape of the erythrocyte membrane (PubMed:35835865). Glycophorin A is the major intrinsic membrane protein of the erythrocyte. The N-terminal glycosylated segment, which lies outside the erythrocyte membrane, has MN blood group receptors. Appears to be important for the function of SLC4A1 and is required for high activity of SLC4A1. May be involved in translocation of SLC4A1 to the plasma membrane.

Cellular Location
Cell membrane; Single-pass type I membrane protein Note=Appears to be colocalized with SLC4A1


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Glycophorin A / CD235a (Erythrocyte Marker) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Glycophorin A / CD235a (Erythrocyte Marker) Antibody - With BSA and Azide - Images



Formalin-fixed, paraffin-embedded human Placenta stained with Glycophorin A Monoclonal Antibody (SPM599)

Glycophorin A / CD235a (Erythrocyte Marker) Antibody - With BSA and Azide - Background

Recognizes a sialoglycoprotein of 39kDa, identified as glycophorin A (GPA). It is present on red blood cells (RBC) and erythroid precursor cells. It has been shown that glycophorin acts as the receptor for Sandei virus and parvovirus. Glycophorins A (GPA) and B (GPB), which are single, trans-membrane sialoglycoproteins. GPA is the carrier of blood group M and N specificities, while GPB accounts for S and U specificities. GPA and GPB provide the cells with a large mucin like surface and it has been suggested this provides a barrier to cell fusion, so minimizing aggregation between red blood cells in the circulation.

Glycophorin A / CD235a (Erythrocyte Marker) Antibody - With BSA and Azide - References

Andersson, L.C., et al. 1979. Glycophorin A as a cell surface marker of early erythroid differentiation in acute leukemia. Int. J. Cancer 23: 717-720. | Liszka, K., et al., 1983. Glycophorin A expression in malignant hematopoiesis. Am. J. Hematol. 15: 219-226. |