10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CD25 / IL2RA (Activated Lymphocyte Marker) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone 143-13]
Catalog # AH11574

Specification

CD25 / IL2RA (Activated Lymphocyte Marker) Antibody - With BSA and Azide - Product Information

Application	**,3,4,**
Primary Accession	P01589
Other Accession	3559, 231367
Reactivity	**Human**
Host	**Mouse**
Clonality	**Monoclonal**
Isotype	**Mouse / IgG1, kappa**
Calculated MW	**55kDa KDa**

CD25 / IL2RA (Activated Lymphocyte Marker) Antibody - With BSA and Azide - Additional Information

Gene ID 3559

Other Names
Interleukin-2 receptor subunit alpha, IL-2 receptor subunit alpha, IL-2-RA, IL-2R subunit alpha, IL2-RA, TAC antigen, p55, CD25, IL2RA

Storage
Store at 2 to 8°C.Antibody is stable for 24 months.

Precautions
CD25 / IL2RA (Activated Lymphocyte Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

CD25 / IL2RA (Activated Lymphocyte Marker) Antibody - With BSA and Azide - Protein Information

Name IL2RA

Function
Receptor for interleukin-2. The receptor is involved in the regulation of immune tolerance by controlling regulatory T cells (TREGs) activity. TREGs suppress the activation and expansion of autoreactive T-cells.

Cellular Location
Membrane; Single-pass type I membrane protein.

CD25 / IL2RA (Activated Lymphocyte Marker) Antibody - With BSA and Azide - Protocols

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CD25 / IL2RA (Activated Lymphocyte Marker) Antibody - With BSA and Azide - Images

CD25 / IL2RA (Activated Lymphocyte Marker) Antibody - With BSA and Azide - Background

Recognizes a protein of 55kDa, identified as CD25 (Workshop IV; Code A27). CD25 is expressed on activated T- and B-cells and activated monocytes/macrophages. With respect to lymphomas, CD25 is present on malignant cells of Hodgkinā€™s disease, HTLV-1+ adult T-cell leukemia, cutaneous T-cell lymphoma, and hair cell leukemia. Increased levels of soluble CD25 are observed in the leukemias/lymphomas and inflammatory/ autoimmune diseases. CD25 molecule alone appears to function as a low affinity receptor and associates with CD122 (IL-2R ļ€ € chain, p75) and CD132 (common ļ€ € chain) to form the high affinity IL-2 receptor complex. CD25 antibodies detect three epitope regions, A, B and C. This MAb recognizes the epitope B, which is located at residue 3-104 of CD25 and can effectively block IL-2 binding to CD25.

CD25 / IL2RA (Activated Lymphocyte Marker) Antibody - With BSA and Azide - References

Knapp W. et al. Leucocyte typing IV, p. 408- 411 and p. 1080, Oxford University Press, Oxford 1989