

**GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide**  
**Rat Monoclonal Antibody [Clone HSP90B1/1192 ]**  
**Catalog # AH12459**

**Specification**

**GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide - Product Information**

|                   |   |
|-------------------|---|
| Application       | ,1,14,3,4,                                    |
| Primary Accession | <a href="#">P14625</a>                        |
| Other Accession   | <a href="#">7184</a> , <a href="#">192374</a> |
| Reactivity        | Human   |
| Host              | Rat   |
| Clonality         | Monoclonal                                    |
| Isotype           | Rat / IgG2a, kappa                            |
| Calculated MW     | 94kDa KDa                                     |

**GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide - Additional Information**

**Gene ID** 7184

**Other Names**

Endoplasmic, 94 kDa glucose-regulated protein, GRP-94, Heat shock protein 90 kDa beta member 1, Tumor rejection antigen 1, gp96 homolog, HSP90B1, GRP94, TRA1

**Storage**

Store at 2 to 8°C. Antibody is stable for 24 months.

**Precautions**

GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

**GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide - Protein Information**

**Name** HSP90B1 ([HGNC:12028](#))

**Synonyms** GRP94, TRA1

**Function**

Molecular chaperone that functions in the processing and transport of secreted proteins (By similarity). When associated with CNPY3, required for proper folding of Toll-like receptors (By similarity). Functions in endoplasmic reticulum associated degradation (ERAD) (PubMed: [18264092](http://www.uniprot.org/citations/18264092)). Has ATPase activity (By similarity). May participate in the unfolding of cytosolic leaderless cargos (lacking the secretion signal sequence) such as the interleukin 1/IL-1 to facilitate their translocation into the ERGIC (endoplasmic reticulum- Golgi intermediate compartment) and secretion; the translocation process is mediated by the cargo receptor TMED10 (PubMed: [a](#)

href="http://www.uniprot.org/citations/32272059" target="\_blank">32272059</a>).

#### **Cellular Location**


Endoplasmic reticulum lumen. Sarcoplasmic reticulum lumen {ECO:0000250|UniProtKB:P41148}.  
Melanosome Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

#### **GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide - Images**



Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with GRP94 Monoclonal Antibody (HSP90B1/1192).

#### **GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide - Background**

Recognizes a protein of 94kDa, which is identified as the glucose-regulated protein 94 (grp94) and also tumor rejection antigen (gp96). Grp94 shows a high degree of sequence homology with the heat shock protein 90 (hsp90). This MAb is highly specific to grp94 and shows minimal cross-reaction with other members of the HSP90 family. Grp s are a class of proteins unresponsive to heat shock and are induced by glucose deprivation. Grp94 has been briefly studied as a prognostic factor in breast cancer.

#### **GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide - References**

Sorger, P.K. et al. J. Mol. Biol. 194: 341-344 (1987). | Tandon, A.K. et.al. Breast Cancer Res. and

Treat. 16: 146 (1990). |