

Lewis A (Blood Group Antigen) Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone 7LE]
Catalog # AH12865

Specification

Lewis A (Blood Group Antigen) Antibody - With BSA and Azide - Product Information

Application	,2,3,4,
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Calculated MW	Multiple KDa

Lewis A (Blood Group Antigen) Antibody - With BSA and Azide - Additional Information

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

Lewis A (Blood Group Antigen) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.


Lewis A (Blood Group Antigen) Antibody - With BSA and Azide - Protein Information


Lewis A (Blood Group Antigen) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Lewis A (Blood Group Antigen) Antibody - With BSA and Azide - Images





Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with Lewis A Monoclonal Antibody (7LE).

Lewis A (Blood Group Antigen) Antibody - With BSA and Azide - Background

Recognizes a carbohydrate determinant of Gal 1-3(Fuc 1-4) GlcNAc which is blood group antigen Lewis A. It is present primarily on epithelial cells such as colon and kidneys. In the tumors and dedifferentiated tissues, decrease of Lewis A antigen was observed. Lewis A (type 1 chain) is expressed in colonic epithelial cells and may be useful for detection of gastrointestinal tumors, pancreatic cancer, and colorectal tumors. Blood group related antigens represent a group of carbohydrate determinants carried on both glycolipids and glycoproteins. They are usually mucin-type, and are detected on erythrocytes, certain epithelial cells, and in secretions of certain individuals. Sixteen genetically and biosynthetically distinct but inter-related specificities belong to this group of antigens, including A, B, H, Lewis A, Lewis B, Lewis X, Lewis Y, and precursor type 1 chain antigens.

Lewis A (Blood Group Antigen) Antibody - With BSA and Azide - References

Blood transfusion and immunohaematology, Ph. Rouger, D Anstee and Ch. Salmon (Eds). Arnette, France 30 (5): 353-720 (1987). | Cancer Epidemiology, Biomarkers Prevention 1, 199-205 (1992). | Gastroenterology, 102 (2), 424-430 (1992)