

NPAS1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10040**Specification**

NPAS1 antibody - C-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q99742
Other Accession	Q99742 , NP_002508 , NM_002517
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	63 kDa KDa

NPAS1 antibody - C-terminal region - Additional Information**Gene ID 4861**

Alias Symbol **MOP5, PASD5, bHLHe11**

Other Names

Neuronal PAS domain-containing protein 1, Neuronal PAS1, Basic-helix-loop-helix-PAS protein MOP5, Class E basic helix-loop-helix protein 11, bHLHe11, Member of PAS protein 5, PAS domain-containing protein 5, NPAS1, BHLHE11, MOP5, PASD5

Target/Specificity

NPAS1 is a member of the basic helix-loop-helix (bHLH)-PAS family of transcription factors. Studies of a related mouse gene suggest that it functions in neurons. The exact function of this gene is unclear, but it may play protective or modulatory roles during late embryogenesis and postnatal development.

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-NPAS1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

NPAS1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NPAS1 antibody - C-terminal region - Protein Information

Name NPAS1

Synonyms BHLHE11, MOP5, PASD5

Function

May control regulatory pathways relevant to schizophrenia and to psychotic illness. May play a role in late central nervous system development by modulating EPO expression in response to cellular oxygen level (By similarity). Forms a heterodimer that binds core DNA sequence 5'-TACGTG-3' within the hypoxia response element (HRE) leading to transcriptional repression on its target gene TH (By similarity).


Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00981}.


NPAS1 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NPAS1 antibody - C-terminal region - Images

NPAS1 antibody - C-terminal region (AI10040) in Human Stomach cells using Immunohistochemistry
Human Stomach



NPAS1 antibody - C-terminal region (AI10040) in Human Jurkat cells using Western Blot
WB Suggested Anti-NPAS1 Antibody Titration: 0.2-1 µg/ml

Positive Control: Jurkat cell lysate

NPAS1 is supported by BioGPS gene expression data to be expressed in Jurkat

NPAS1 antibody - C-terminal region - Background

This is a rabbit polyclonal antibody against NPAS1. It was validated on Western Blot and immunohistochemistry by Abgent. At Abgent we manufacture rabbit polyclonal antibodies on a large scale (200-1000 products/month) of high throughput manner. Our antibodies are peptide based and protein family oriented. We usually provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).